




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 16/00		A2	(11) International Publication Number: WO 98/50433
			(43) International Publication Date: 12 November 1998 (12.11.98)
(21) International Application Number: PCT/US98/09160 (22) International Filing Date: 5 May 1998 (05.05.98) (30) Priority Data: 08/851,362 5 May 1997 (05.05.97) US (71) Applicant: ABGENIX, INC. [US/US]; 7601 Dumbarton Circle, Fremont, CA 94555 (US). (72) Inventors: JAKOBOVITS, Aya; 2021 Monterey Avenue, Menlo Park, CA 94025 (US). YANG, Xiao-Dong; 2833 Bryant Street, Palo Alto, CA 94306 (US). GALLO, Michael; 1376 Klamath Drive, San Jose, CA 95130 (US). JIA, Xiao-Chi; 64 Burbank Avenue, San Mateo, CA 94403 (US). (74) Agents: HALEY, James, F. et al.; Fish & Neave, 1251 Avenue of the Americas, New York, NY 10020 (US).			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report.

(54) Title: MAN MONOCLONAL ANTIBODIES TO EPIDERMAL GROWTH FACTOR RECEPTOR



(57) Abstract

In accordance with the present invention, there are provided fully human monoclonal antibodies against human epidermal growth factor receptor (EGF-r). Nucleotide sequences encoding and amino acid sequences comprising heavy and light chain immunoglobulin molecules, particularly sequences corresponding to contiguous heavy and light chain sequences from CDR1 through CDR3, are provided. Hybridomas expressing such immunoglobulin molecules and monoclonal antibodies are also provided.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

HUMAN MONOCLONAL ANTIBODIES TO EPIDERMAL GROWTH FACTOR
RECEPTOR

BACKGROUND OF THE INVENTION

1. Summary of the Invention

5 In accordance with the present invention, there are provided fully human contiguous heavy and light chain sequences spanning the complementarity determining regions monoclonal antibodies against human epidermal growth factor receptor (EGF-r). Nucleotide
10 sequences encoding and amino acid sequences comprising heavy and light chain immunoglobulin molecules, particularly sequences corresponding to (CDR's), specifically from CDR1 through CDR3, are provided. Hybridomas expressing such immunoglobulin molecules and
15 monoclonal antibodies are also provided.

2. Background of the Technology

EGF-r has been demonstrated to be overexpressed on many types of human solid tumors. Mendelsohn *Cancer Cells* 7:359 (1989), Mendelsohn *Cancer Biology* 1:339-344
20 (1990), Modjtahedi and Dean *Int'l J. Oncology* 4:277-296

-2-

(1994). For example, EGF-r overexpression has been observed in certain lung, breast, colon, gastric, brain, bladder, head and neck, ovarian, and prostate carcinomas. Modjtahedi and Dean *Int'l J. Oncology* 4:277-296 (1994). Both epidermal growth factor (EGF) and transforming growth factor-alpha (TGF- α) have been demonstrated to bind to EGF-r and to lead to cellular proliferation and tumor growth.

Thus, certain groups have proposed that antibodies against EGF, TGF- α , and EGF-r may be useful in the therapy of tumors expressing or overexpressing EGF-r. Mendelsohn *Cancer Cells* 7:359 (1989), Mendelsohn *Cancer Biology* 1:339-344 (1990), Modjtahedi and Dean *Int'l J. Oncology* 4:277-296 (1994), Tosi et al. *Int'l J. Cancer* 62:643-650 (1995). Indeed, it has been demonstrated that anti-EGF-r antibodies while blocking EGF and TGF- α binding to the receptor appear to inhibit tumor cell proliferation. At the same time, however, anti-EGF-r antibodies have not appeared to inhibit EGF and TGF- α independent cell growth. Modjtahedi and Dean *Int'l J. Oncology* 4:277-296 (1994).

In view of these findings, a number of murine and rat monoclonal antibodies against EGF-r have been developed and tested for their ability inhibit the growth of tumor cells in vitro and in vivo. Modjtahedi and Dean *Int'l J. Oncology* 4:277-296 (1994). The antibody that has apparently advanced the farthest in the clinic is a chimeric antibody, designated C225, which has a murine variable region and a human IgG1 constant region. Modjtahedi and Dean *Int'l J. Oncology* 4:277-296 (1994). The murine antibody, designated 225,

upon which the C225 antibody is based, was developed by University of California and Rorer. See U.S. Patent No. 4,943,533 and European Patent No. 359,282, the disclosures of which are hereby incorporated by
5 reference. The C225 antibody was demonstrated to inhibit EGF-mediated tumor cell growth in vitro and inhibit human tumor formation in vivo in nude mice. The antibody, moreover, appeared to act in synergy with certain chemotherapeutic agents to eradicate human
10 tumors in vivo in xenograft mouse models. Modjtahedi and Dean *Int'l J. Oncology* 4:277-296 (1994).

ImClone has been conducting human clinical trials using the anti-EGF-r antibody designated C225. Phase I and Phase I/II clinical trials in patients with head
15 and neck, prostate, and lung carcinomas apparently have been, or are currently being, conducted with C225. In Phase I clinical trials, no toxicity was detected with multiple injections and with doses of up to perhaps 400 mg/m², even in cases involving immunocompromised
20 patients. Such studies were conducted as dose escalation studies comprising 5 doses of from about 5 to about 200 mg/m² and were performed in combination with chemotherapy (i.e., doxorubicin, adriamycin, taxol, and cisplatin). In addition to the apparent
25 safety data that has been generated in these studies, preliminary results from the studies appear to indicate some evidence of tumor shrinkage in 80% of patients having prostate cancer.

Each of these above-mentioned antibodies, however,
30 possess murine or rat variable and/or constant regions. The presence of such murine or rat derived proteins can lead to the rapid clearance of the antibodies or can

- 4 -

lead to the generation of an immune response against the antibody by a patient. In order to avoid the utilization of murine or rat derived antibodies, it has been postulated that one could introduce human antibody
5 function into a rodent so that the rodent would produce fully human antibodies.

The ability to clone and reconstruct megabase-sized human loci in YACs and to introduce them into the mouse germline provides a powerful approach to
10 elucidating the functional components of very large or crudely mapped loci as well as generating useful models of human disease. Furthermore, the utilization of such technology for substitution of mouse loci with their human equivalents could provide unique insights into
15 the expression and regulation of human gene products during development, their communication with other systems, and their involvement in disease induction and progression.

An important practical application of such a
20 strategy is the "humanization" of the mouse humoral immune system. Introduction of human immunoglobulin (Ig) loci into mice in which the endogenous Ig genes have been inactivated offers the opportunity to study the mechanisms underlying programmed expression and
25 assembly of antibodies as well as their role in B-cell development. Furthermore, such a strategy could provide an ideal source for production of fully human monoclonal antibodies (Mabs) - an important milestone towards fulfilling the promise of antibody therapy in
30 human disease. Fully human antibodies are expected to minimize the immunogenic and allergic responses intrinsic to mouse or mouse-derivatized Mabs and thus

-5-

to increase the efficacy and safety of the administered antibodies. The use of fully human antibodies can be expected to provide a substantial advantage in the treatment of chronic and recurring human diseases, such as inflammation, autoimmunity, and cancer, which require repeated antibody administrations.

One approach towards this goal was to engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig loci in anticipation that such mice would produce a large repertoire of human antibodies in the absence of mouse antibodies. Large human Ig fragments would preserve the large variable gene diversity as well as the proper regulation of antibody production and expression. By exploiting the mouse machinery for antibody diversification and selection and the lack of immunological tolerance to human proteins, the reproduced human antibody repertoire in these mouse strains should yield high affinity antibodies against any antigen of interest, including human antigens. Using the hybridoma technology, antigen-specific human Mabs with the desired specificity could be readily produced and selected.

This general strategy was demonstrated in connection with our generation of the first XenoMouse™ strains as published in 1994. See Green et al. *Nature Genetics* 7:13-21 (1994). The XenoMouse™ strains were engineered with yeast artificial chromosomes (YACs) containing 245 kb and 190 kb-sized germline configuration fragments of the human heavy chain locus and kappa light chain locus, respectively, which contained core variable and constant region sequences.

-6-

Id. The human Ig containing YACs proved to be compatible with the mouse system for both rearrangement and expression of antibodies and were capable of substituting for the inactivated mouse Ig genes. This was demonstrated by their ability to induce B-cell development, to produce an adult-like human repertoire of fully human antibodies, and to generate antigen-specific human Mabs. These results also suggested that introduction of larger portions of the human Ig loci containing greater numbers of V genes, additional regulatory elements, and human Ig constant regions might recapitulate substantially the full repertoire that is characteristic of the human humoral response to infection and immunization. The work of Green et al. was recently extended to the introduction of greater than approximately 80% of the human antibody repertoire through introduction of megabase sized, germline configuration YAC fragments of the human heavy chain loci and kappa light chain loci, respectively. See Mendez et al. *Nature Genetics* 15:146-156 (1997) and U.S. Patent Application Serial No. 08/759,620, filed December 3, 1996, the disclosures of which are hereby incorporated by reference.

Such approach is further discussed and delineated in U.S. Patent Application Serial Nos. 07/466,008, filed January 12, 1990, 07/610,515, filed November 8, 1990, 07/919,297, filed July 24, 1992, 07/922,649, filed July 30, 1992, filed 08/031,801, filed March 15, 1993, 08/112,848, filed August 27, 1993, 08/234,145, filed April 28, 1994, 08/376,279, filed January 20, 1995, 08/430, 938, April 27, 1995, 08/464,584, filed June 5, 1995, 08/464,582, filed June 5, 1995,

-7-

08/463,191, filed June 5, 1995, 08/462,837, filed June 5, 1995, 08/486,853, filed June 5, 1995, 08/486,857, filed June 5, 1995, 08/486,859, filed June 5, 1995, 08/462,513, filed June 5, 1995, 08/724,752, filed
5 October 2, 1996, and 08/759,620, filed December 3, 1996. See also Mendez et al. *Nature Genetics* 15:146-156 (1997). See also European Patent No., EP 0 463 151 B1, grant published June 12, 1996, International Patent Application No., WO 94/02602, published February 3,
10 1994, International Patent Application No., WO 96/34096, published October 31, 1996, and PCT Application No. PCT/US96/05928, filed April 29, 1996. The disclosures of each of the above-cited patents, applications, and references are hereby incorporated by
15 reference in their entirety.

In an alternative approach, others, including GenPharm International, Inc., have utilized a "minilocus" approach. In the minilocus approach, an exogenous Ig locus is mimicked through the inclusion of
20 pieces (individual genes) from the Ig locus. Thus, one or more V_H genes, one or more D_H genes, one or more J_H genes, a mu constant region, and a second constant region (preferably a gamma constant region) are formed into a construct for insertion into an animal. This
25 approach is described in U.S. Patent No. 5,545,807 to Surani et al. and U.S. Patent Nos. 5,545,806 and 5,625,825, both to Lonberg and Kay, and GenPharm International U.S. Patent Application Serial Nos. 07/574,748, filed August 29, 1990, 07/575,962, filed
30 August 31, 1990, 07/810,279, filed December 17, 1991, 07/853,408, filed March 18, 1992, 07/904,068, filed June 23, 1992, 07/990,860, filed December 16, 1992,

08/053,131, filed April 26, 1993, 08/096,762, filed
July 22, 1993, 08/155,301, filed November 18, 1993,
08/161,739, filed December 3, 1993, 08/165,699, filed
December 10, 1993, 08/209,741, filed March 9, 1994, the
5 disclosures of which are hereby incorporated by
reference. See also International Patent Application
Nos. WO 94/25585, published November 10, 1994, WO
93/12227, published June 24, 1993, WO 92/22645,
published December 23, 1992, WO 92/03918, published
10 March 19, 1992, the disclosures of which are hereby
incorporated by reference in their entirety. See
further Taylor et al., 1992, Chen et al., 1993,
Tuailon et al., 1993, Choi et al., 1993, Lonberg et
al., (1994), Taylor et al., (1994), and Tuailon et
15 al., (1995), the disclosures of which are hereby
incorporated by reference in their entirety.

The inventors of Surani et al., cited above and
assigned to the Medical Research Counsel (the "MRC"),
produced a transgenic mouse possessing an Ig locus
20 through use of the minilocus approach. The inventors
on the GenPharm International work, cited above,
Lonberg and Kay, following the lead of the present
inventors, proposed inactivation of the endogenous
mouse Ig locus coupled with substantial duplication of
25 the Surani et al. work.

An advantage of the minilocus approach is the
rapidity with which constructs including portions of
the Ig locus can be generated and introduced into
animals. Commensurately, however, a significant
30 disadvantage of the minilocus approach is that, in
theory, insufficient diversity is introduced through

-9-

the inclusion of small numbers of V, D, and J genes. Indeed, the published work appears to support this concern. B-cell development and antibody production of animals produced through use of the minilocus approach appear stunted. Therefore, research surrounding the present invention has consistently been directed towards the introduction of large portions of the Ig locus in order to achieve greater diversity and in an effort to reconstitute the immune repertoire of the animals.

Human anti-mouse antibody (HAMA) responses have led the industry to prepare chimeric or otherwise humanized antibodies. While the C225 antibody is a chimeric antibody, having a human constant region and a murine variable region, it is expected that certain human anti-chimeric antibody (HACA) responses will be observed, particularly in chronic or multi-dose utilizations of the antibody.

Thus, it would be desirable to provide fully human antibodies against EGF-r that possess similar or enhanced activities as compared to C225 in order to vitiate concerns and/or effects of HAMA or HACA response.

BRIEF DESCRIPTION OF THE DRAWING FIGURES

Figure 1 is an amino acid sequence of a heavy chain immunoglobulin molecule that is secreted by the hybridoma E1.1. Differences between the sequence encoded by heavy chain variable gene 4-31 and the sequence of the E1.1 secreted heavy chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and

-10-

CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 2 is a nucleotide sequence of the cDNA encoding the heavy chain immunoglobulin molecule of Figure 1 that was cloned out of the hybridoma E1.1.

Figure 3 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E1.1. Differences between the sequence encoded by light chain variable gene 018 and the sequence of the E1.1 secreted light chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 4 is a nucleotide sequence of the cDNA encoding the kappa light chain immunoglobulin molecule of Figure 3 that was cloned out of the hybridoma E1.1.

Figure 5 is an amino acid sequence of a heavy chain immunoglobulin molecule that is secreted by the hybridoma E2.4. Differences between the sequence encoded by heavy chain variable gene 4-31 and the sequence of the E2.4 secreted heavy chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 6 is a nucleotide sequence of the cDNA encoding the heavy chain immunoglobulin molecule of Figure 5 that was cloned out of the hybridoma E2.4.

Figure 7 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E2.4. Differences between the sequence

encoded by light chain variable gene 018 and the sequence of the E2.4 secreted light chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 8 is a nucleotide sequence of the cDNA encoding the kappa light chain immunoglobulin molecule of Figure 7 that was cloned out of the hybridoma E2.4.

Figure 9 is an amino acid sequence of a heavy chain immunoglobulin molecule that is secreted by the hybridoma E2.5. Differences between the sequence encoded by heavy chain variable gene 4-31 and the sequence of the E2.5 secreted heavy chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 10 is a nucleotide sequence of the cDNA encoding the heavy chain immunoglobulin molecule of Figure 9 that was cloned out of the hybridoma E2.5.

Figure 11 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E2.5. Differences between the sequence encoded by light chain variable gene 018 and the sequence of the E2.5 secreted light chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 12 is a nucleotide sequence of the cDNA encoding the kappa light chain immunoglobulin molecule

- 12 -

of Figure 11 that was cloned out of the hybridoma E2.5.

Figure 13 is an amino acid sequence of a heavy chain immunoglobulin molecule that is secreted by the hybridoma E6.2. Differences between the sequence encoded by heavy chain variable gene 4-31 and the sequence of the E6.2 secreted heavy chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 14 is a nucleotide sequence of the cDNA encoding the heavy chain immunoglobulin molecule of Figure 13 that was cloned out of the hybridoma E6.2.

Figure 15 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E6.2. Differences between the sequence encoded by light chain variable gene 018 and the sequence of the E6.2 secreted light chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 16 is a nucleotide sequence of the cDNA encoding the kappa light chain immunoglobulin molecule of Figure 15 that was cloned out of the hybridoma E6.2.

Figure 17 is an amino acid sequence of a heavy chain immunoglobulin molecule that is secreted by the hybridoma E6.4. Differences between the sequence encoded by heavy chain variable gene 4-31 and the sequence of the E6.4 secreted heavy chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and

- 13 -

CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 18 is a nucleotide sequence of the cDNA encoding the heavy chain immunoglobulin molecule of Figure 17 that was cloned out of the hybridoma E6.2.

Figure 19 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E6.4. Differences between the sequence encoded by light chain variable gene 018 and the sequence of the E6.4 secreted light chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 20 is a nucleotide sequence of the cDNA encoding the kappa light chain immunoglobulin molecule of Figure 19 that was cloned out of the hybridoma E6.4.

Figure 21 is an amino acid sequence of a heavy chain immunoglobulin molecule that is secreted by the hybridoma E2.11. Differences between the sequence encoded by heavy chain variable gene 4-61 and the sequence of the E2.11 secreted heavy chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 22 is a nucleotide sequence of the cDNA encoding the heavy chain immunoglobulin molecule of Figure 21 that was cloned out of the hybridoma E2.11.

Figure 23 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E2.11. Differences between the sequence

encoded by light chain variable gene 018 and the sequence of the E2.11 secreted light chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 24 is a nucleotide sequence of the cDNA encoding the kappa light chain immunoglobulin molecule of Figure 23 that was cloned out of the hybridoma E2.11.

Figure 25 is an amino acid sequence of a heavy chain immunoglobulin molecule that is secreted by the hybridoma E6.3. Differences between the sequence encoded by heavy chain variable gene 4-61 and the sequence of the E6.3 secreted heavy chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 26 is a nucleotide sequence of the cDNA encoding the heavy chain immunoglobulin molecule of Figure 25 that was cloned out of the hybridoma E6.3.

Figure 27 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E6.3. Differences between the sequence encoded by light chain variable gene 018 and the sequence of the E6.3 secreted light chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 28 is a nucleotide sequence of the cDNA

encoding the kappa light chain immunoglobulin molecule of Figure 27 that was cloned out of the hybridoma E6.3.

Figure 29 is an amino acid sequence of a heavy chain immunoglobulin molecule that is secreted by the hybridoma E7.6.3. Differences between the sequence encoded by heavy chain variable gene 4-61 and the sequence of the E7.6.3 secreted heavy chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 30 is a nucleotide sequence of the cDNA encoding the heavy chain immunoglobulin molecule of Figure 29 that was cloned out of the hybridoma E7.6.3.

Figure 31 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E7.6.3. Differences between the sequence encoded by light chain variable gene 018 and the sequence of the E7.6.3 secreted light chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 32 is a nucleotide sequence of the cDNA encoding the kappa light chain immunoglobulin molecule of Figure 31 that was cloned out of the hybridoma E7.6.3.

Figure 33 provides a comparison of specific anti-EGF-r antibody heavy chain amino acid sequence comparisons with the amino acid sequence of the particular V_H gene which encodes the heavy chain of the particular antibody.

Figure 34 provides a comparison of specific anti-EGF-r antibody light chain amino acid sequence comparisons with the amino acid sequence of the particular Vk gene which encodes the light chain of the particular antibody.

Figure 35 shows blockage EGF binding to human epidermoid carcinoma A431 cells by human anti-EGF-r antibodies in vitro, where (□) depicts the results achieved by an anti-EGF-r antibody in accordance with the invention, (●) depicts the results achieved by the murine monoclonal antibody 225, and (▲) depicts the results achieved by a control, nonspecific, human IgG2 antibody.

Figure 36 shows inhibition of EGF binding to human epidermoid carcinoma A431 cells by human anti-EGF-r antibodies in vitro, where (□) depicts the results achieved by the murine monoclonal antibody 225, (○) depicts the results achieved by the murine monoclonal antibody 528, (▼) depicts the results achieved using the E1.1 antibody in accordance with the invention, (▲) depicts the results achieved using the E2.4 antibody in accordance with the invention, (▸) depicts the results achieved using the E2.5 antibody in accordance with the invention, (◄) depicts the results achieved using the E2.6 antibody in accordance with the invention, (◆) depicts the results achieved using the E2.11 antibody in accordance with the invention, and (⊗) depicts the results achieved using a control, nonspecific human IgG2 antibody.

Figure 37 shows inhibition of TGF- α binding to human epidermoid carcinoma A431 cells by human anti-EGF-r antibodies in vitro, where (□) depicts the

results achieved by the murine monoclonal antibody 225, (◆) depicts the results achieved using the E6.2 antibody in accordance with the invention, (●) depicts the results achieved using the E6.3 antibody in accordance with the invention, (▲) depicts the results achieved using the E7.2 antibody in accordance with the invention, (■) depicts the results achieved using the E7.10 antibody in accordance with the invention, (▼) depicts the results achieved using the E7.6.3, and (⊗) depicts the results achieved using a control, nonspecific human IgG2 antibody.

Figure 38 shows inhibition of EGF binding to human colon carcinoma SW948 cells by human anti-EGF-r antibodies in vitro, where (●) depicts the results achieved by an anti-EGF-r antibody in accordance with the invention, (□) depicts the results achieved by the murine monoclonal antibody 225, and (▲) depicts the results achieved by a control, nonspecific, human IgG2 antibody.

Figure 39 shows that human anti-EGF-r antibodies derived from XenoMouse II strains inhibit growth of SW948 cells in vitro, where (○) depicts the results achieved by an anti-EGF-r antibody in accordance with the invention, (□) depicts the results achieved by the murine monoclonal antibody 225, and (▲) depicts the results achieved by a control, nonspecific, human IgG2 antibody.

Figure 40 shows the inhibition of human epidermoid carcinoma A431 cell growth in nude mice through use of human anti-EGF-r antibodies in accordance with the invention in vivo. In the Figure, (▲) depicts the results achieved with a dosage of 1 mg of a human anti-

EGF-r antibody in accordance with the present invention, (▼) depicts the results achieved with a dosage of 0.2 mg of a human anti-EGF-r antibody in accordance with the present invention, (□) depicts the results achieved by a control, nonspecific, human IgG2 antibody, and (O) depicts the results achieved utilizing phosphate buffered saline as a control.

Figure 41 shows data related to the inhibition of epidermoid carcinoma formation in nude mice through use of human anti-EGF-r antibodies in accordance with the invention in vivo showing tumor incidence at day 19.

Figure 42 shows data related to the inhibition of epidermoid carcinoma formation in nude mice through use of human anti-EGF-r antibodies in accordance with the invention in vivo showing tumor incidence at day 120.

Figure 43 shows data related to the eradication of an established human epidermoid tumor in nude mice through use of human anti-EGF-r antibodies in accordance with the invention in vivo. In the Figure, (▲) depicts the results achieved with multiple doses of 1 mg each of a human anti-EGF-r antibody in accordance with the present invention (E7.6.3), (X) depicts the results achieved with two doses of 125 µg each of doxorubicin, (*) depicts the results achieved with a multiple doses of 1 mg each of a human anti-EGF-r antibody in accordance with the present invention (E7.6.3) in combination with two doses of 125 µg each of doxorubicin, (■) depicts the results achieved by a control, nonspecific, human IgG2 antibody, and (◆) depicts the results achieved utilizing phosphate buffered saline as a control.

Figure 44 shows data related to the eradication of

- 19 -

an established human epidermoid tumor in nude mice through use of human anti-EGF-r antibodies in accordance with the invention in vivo. In the Figure, (◆) depicts the results achieved with multiple doses of 0.5 mg each of a human anti-EGF-r antibody in accordance with the present invention (E2.5), (■) depicts the results achieved with two doses of 125 µg each of doxorubicin, (▲) depicts the results achieved with multiple doses of 0.5 mg each of a human anti-EGF-r antibody in accordance with the present invention (E2.5) in combination with two doses of 125 µg each of doxorubicin, (X) depicts the results achieved utilizing phosphate buffered saline as a control, and (*) depicts the results achieved utilizing a control, nonspecific, human IgG2 antibody at a dose of 1 mg.

SUMMARY OF THE INVENTION

In accordance with a first aspect of the present invention, there is provided an antibody against epidermal growth factor receptor comprising a heavy chain variable region amino acid sequence wherein a portion of the sequence is encoded by a human V_H 4 family gene and any of the mutations thereto represented by the nucleotide sequences shown in Figures 2, 6, 10, 14, 18, 22, 26, and 30. In a preferred embodiment, the heavy chain variable region amino acid sequence comprises an Aspartic Acid amino acid substitution at residue 10.

In accordance with a second aspect of the present invention, there is provided an antibody against epidermal growth factor receptor comprising a heavy chain variable region amino acid sequence wherein a

-20-

portion of the sequence is encoded by a human V_H 4-31 gene and any of the mutations thereto represented by the nucleotide sequences shown in Figures 2, 6, 10, 14, and 18. In a preferred embodiment, the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:23. In a preferred embodiment, the antibody further comprises a light chain variable region comprising the sequence represented by SEQ ID NO:24. In a preferred embodiment, the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:25. In a preferred embodiment, the antibody further comprises a light chain variable region comprising the sequence represented by SEQ ID NO:26. In a preferred embodiment, the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:27. In a preferred embodiment, the antibody further comprises a light chain variable region comprising the sequence represented by SEQ ID NO:28. In a preferred embodiment, the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:29. In a preferred embodiment, the antibody further comprises a light chain variable region comprising the sequence represented by SEQ ID NO:30. In a preferred embodiment, the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:31. In a preferred embodiment, the antibody further comprises a light chain variable region comprising the sequence represented by SEQ ID NO:32.

In accordance with the third aspect of the present

-21-

invention, there is provided an antibody against epidermal growth factor receptor comprising a heavy chain variable region amino acid sequence wherein a portion of the sequence is encoded by a human V_H 4-61 gene and any of the mutations thereto represented by the nucleotide sequences shown in Figures 22, 26, and 30. In a preferred embodiment, the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:33. In a preferred embodiment, the antibody further comprises a light chain variable region comprising the sequence represented by SEQ ID NO:34. In a preferred embodiment, the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:35. In a preferred embodiment, the antibody further comprises a light chain variable region comprising the sequence represented by SEQ ID NO:36. In a preferred embodiment, the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:37. In a preferred embodiment, the antibody further comprises a light chain variable region comprising the sequence represented by SEQ ID NO:38.

In accordance with a fourth aspect of the present invention, there is provided an antibody against epidermal growth factor receptor comprising a light chain variable region amino acid sequence wherein a portion of the sequence is encoded by a human V_K I family gene and any of the mutations thereto represented by the nucleotide sequences shown in Figures 4, 8, 12, 16, 20, 24, 28, and 32. In a preferred embodiment, the light chain variable region

- 22 -

comprises the sequence represented by SEQ ID NO:24. In a preferred embodiment, the light chain variable region comprises the sequence represented by SEQ ID NO:26. In a preferred embodiment, the light chain variable region
5 comprises the sequence represented by SEQ ID NO:28. In a preferred embodiment, the light chain variable region comprises the sequence represented by SEQ ID NO:30. In a preferred embodiment, the light chain variable region comprises the sequence represented by SEQ ID NO:32. In
10 a preferred embodiment, the light chain variable region comprises the sequence represented by SEQ ID NO:34. In a preferred embodiment, the light chain variable region comprises the sequence represented by SEQ ID NO:36. In a preferred embodiment, the light chain variable region
15 comprises the sequence represented by SEQ ID NO:38.

In accordance with a fifth aspect of the present invention, there is provided an antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence
20 from CDR1 through CDR3 as represented in SEQ ID NO:23. In a preferred embodiment, the antibody further comprises a light chain variable region comprising the sequence represented by SEQ ID NO:24.

In accordance with a sixth aspect of the present
25 invention, there is provided an antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:25. In a preferred embodiment, the antibody further
30 comprises a light chain variable region comprising the sequence represented by SEQ ID NO:26.

In accordance with a seventh aspect of the present

invention, there is provided an antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:27.

- 5 In a preferred embodiment, the antibody further comprises a light chain variable region comprising the sequence represented by SEQ ID NO:28.

- In accordance with a eighth aspect of the present invention, there is provided an antibody against
10 epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:29. In a preferred embodiment, the antibody further comprises a light chain variable region comprising the
15 sequence represented by SEQ ID NO:30.

- In accordance with a ninth aspect of the present invention, there is provided an antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence
20 from CDR1 through CDR3 as represented in SEQ ID NO:31. In a preferred embodiment, the antibody further comprises a light chain variable region comprising the sequence represented by SEQ ID NO:32.

- In accordance with a tenth aspect of the present
25 invention, there is provided an antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:33. In a preferred embodiment, the antibody further
30 comprises a light chain variable region comprising the sequence represented by SEQ ID NO:34.

In accordance with an eleventh aspect of the

-24-

present invention, there is provided an antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:35. In a preferred embodiment, the antibody further comprises a light chain variable region comprising the sequence represented by SEQ ID NO:36.

In accordance with a twelfth aspect of the present invention, there is provided an antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:37. In a preferred embodiment, the antibody further comprises a light chain variable region comprising the sequence represented by SEQ ID NO:38.

In accordance with a thirteenth aspect of the present invention, there is provided, in a method for treating a solid tumor with an antibody against epidermal growth factor receptor, the improvement comprising administering to a patient having a solid tumor one of the foregoing antibodies of the invention.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

In accordance with the present invention, there are provided fully human monoclonal antibodies against human epidermal growth factor receptor (EGF-r). Nucleotide sequences encoding and amino acid sequences comprising heavy and light chain immunoglobulin molecules, particularly sequences corresponding to a contiguous heavy and light chain sequences from CDR1 through CDR3, are provided. Hybridomas expressing such immunoglobulin molecules and monoclonal antibodies are

also provided.

Definitions

Unless otherwise defined, scientific and technical
5 terms used in connection with the present invention
shall have the meanings that are commonly understood by
those of ordinary skill in the art. Further, unless
otherwise required by context, singular terms shall
include pluralities and plural terms shall include the
10 singular. Generally, nomenclatures utilized in
connection with, and techniques of, cell and tissue
culture, molecular biology, and protein and oligo- or
polynucleotide chemistry and hybridization described
herein are those well known and commonly used in the
15 art. Standard techniques are used for recombinant DNA,
oligonucleotide synthesis, and tissue culture and
transformation (e.g., electroporation, lipofection).
Enzymatic reactions and purification techniques are
performed according to manufacturer's specifications or
20 as commonly accomplished in the art or as described
herein. The foregoing techniques and procedures are
generally performed according to conventional methods
well known in the art and as described in various
general and more specific references that are cited and
25 discussed throughout the present specification. See
e.g., Sambrook et al. Molecular Cloning: A Laboratory
Manual (2d ed., Cold Spring Harbor Laboratory Press,
Cold Spring Harbor, N.Y. (1989)), which is incorporated
herein by reference. The nomenclatures utilized in
30 connection with, and the laboratory procedures and
techniques of, analytical chemistry, synthetic organic
chemistry, and medicinal and pharmaceutical chemistry

-26-

described herein are those well known and commonly used in the art. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment
5 of patients.

As utilized in accordance with the present disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

10

The term "isolated polynucleotide" as used herein shall mean a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, which by virtue of its origin the "isolated polynucleotide" (1)
15 is not associated with all or a portion of a polynucleotide in which the "isolated polynucleotide" is found in nature, (2) is operably linked to a polynucleotide which it is not linked to in nature, or (3) does not occur in nature as part of a larger
20 sequence.

The term "isolated protein" referred to herein means a protein of cDNA, recombinant RNA, or synthetic origin or some combination thereof, which by virtue of its origin, or source of derivation, the "isolated
25 protein" (1) is not associated with proteins found in nature, (2) is free of other proteins from the same source, e.g. free of murine proteins, (3) is expressed by a cell from a different species, or (4) does not occur in nature.

- 27 -

The term "polypeptide" is used herein as a generic term to refer to native protein, fragments, or analogs of a polypeptide sequence. Hence, native protein, fragments, and analogs are species of the polypeptide genus. Preferred polypeptides in accordance with the invention comprise the human heavy chain immunoglobulin molecules represented by Figures 1, 5, 9, 13, 17, 21, 25, and 29 and the human kappa light chain immunoglobulin molecules represented by Figures 3, 7, 11, 15, 19, 23, 27, and 31, as well as antibody molecules formed by combinations comprising the heavy chain immunoglobulin molecules with light chain immunoglobulin molecules, such as the kappa light chain immunoglobulin molecules, and vice versa, as well as fragments and analogs thereof.

The term "naturally-occurring" as used herein as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory or otherwise is naturally-occurring.

The term "operably linked" as used herein refers to positions of components so described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

-28-

The term "control sequence" as used herein refers to polynucleotide sequences which are necessary to effect the expression and processing of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and transcription termination sequence; in eukaryotes, generally, such control sequences include promoters and transcription termination sequence. The term "control sequences" is intended to include, at a minimum, all components whose presence is essential for expression and processing, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences.

The term "polynucleotide" as referred to herein means a polymeric form of nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA.

The term "oligonucleotide" referred to herein includes naturally occurring, and modified nucleotides linked together by naturally occurring, and non-naturally occurring oligonucleotide linkages. Oligonucleotides are a polynucleotide subset generally comprising a length of 200 bases or fewer. Preferably oligonucleotides are 10 to 60 bases in length and most preferably 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length. Oligonucleotides are usually single

-29-

stranded, e.g. for probes; although oligonucleotides may be double stranded, e.g. for use in the construction of a gene mutant. Oligonucleotides of the invention can be either sense or antisense
5 oligonucleotides.

The term "naturally occurring nucleotides" referred to herein includes deoxyribonucleotides and ribonucleotides. The term "modified nucleotides" referred to herein includes nucleotides with modified
10 or substituted sugar groups and the like. The term "oligonucleotide linkages" referred to herein includes oligonucleotides linkages such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate,
15 phoshoraniladate, phosphoroamidate, and the like. See e.g., LaPlanche et al. *Nucl. Acids Res.* 14:9081 (1986); Stec et al. *J. Am. Chem. Soc.* 106:6077 (1984); Stein et al. *Nucl. Acids Res.* 16:3209 (1988); Zon et al. *Anti-Cancer Drug Design* 6:539 (1991); Zon et al.
20 Oligonucleotides and Analogues: A Practical Approach, pp. 87-108 (F. Eckstein, Ed., Oxford University Press, Oxford England (1991)); Stec et al. U.S. Patent No. 5,151,510; Uhlmann and Peyman *Chemical Reviews* 90:543 (1990), the disclosures of which are hereby
25 incorporated by reference. An oligonucleotide can include a label for detection, if desired.

The term "selectively hybridize" referred to herein means to detectably and specifically bind. Polynucleotides, oligonucleotides and fragments thereof

- 30 -

in accordance with the invention selectively hybridize to nucleic acid strands under hybridization and wash conditions that minimize appreciable amounts of detectable binding to nonspecific nucleic acids. High stringency conditions can be used to achieve selective hybridization conditions as known in the art and discussed herein. Generally, the nucleic acid sequence homology between the polynucleotides, oligonucleotides, and fragments of the invention and a nucleic acid sequence of interest will be at least 80%, and more typically with preferably increasing homologies of at least 85%, 90%, 95%, 99%, and 100%. Two amino acid sequences are homologous if there is a partial or complete identity between their sequences. For example, 85% homology means that 85% of the amino acids are identical when the two sequences are aligned for maximum matching. Gaps (in either of the two sequences being matched) are allowed in maximizing matching; gap lengths of 5 or less are preferred with 2 or less being more preferred. Alternatively and preferably, two protein sequences (or polypeptide sequences derived from them of at least 30 amino acids in length) are homologous, as this term is used herein, if they have an alignment score of at more than 5 (in standard deviation units) using the program ALIGN with the mutation data matrix and a gap penalty of 6 or greater. See Dayhoff, M.O., in Atlas of Protein Sequence and Structure, pp. 101-110 (Volume 5, National Biomedical Research Foundation (1972)) and Supplement 2 to this volume, pp. 1-10. The two sequences or parts thereof are more preferably homologous if their amino acids are greater than or equal to 50% identical when optimally

-31-

aligned using the ALIGN program. The term "corresponds to" is used herein to mean that a polynucleotide sequence is homologous (i.e., is identical, not strictly evolutionarily related) to all or a portion of a reference polynucleotide sequence, or that a polypeptide sequence is identical to a reference polypeptide sequence. In contradistinction, the term "complementary to" is used herein to mean that the complementary sequence is homologous to all or a portion of a reference polynucleotide sequence. For illustration, the nucleotide sequence "TATAC" corresponds to a reference sequence "TATAC" and is complementary to a reference sequence "GTATA".

The following terms are used to describe the sequence relationships between two or more polynucleotide or amino acid sequences: "reference sequence", "comparison window", "sequence identity", "percentage of sequence identity", and "substantial identity". A "reference sequence" is a defined sequence used as a basis for a sequence comparison; a reference sequence may be a subset of a larger sequence, for example, as a segment of a full-length cDNA or gene sequence given in a sequence listing or may comprise a complete cDNA or gene sequence. Generally, a reference sequence is at least 18 nucleotides or 6 amino acids in length, frequently at least 24 nucleotides or 8 amino acids in length, and often at least 48 nucleotides or 16 amino acids in length. Since two polynucleotides or amino acid sequences may each (1) comprise a sequence (i.e., a portion of the complete polynucleotide or amino acid

- 32 -

sequence) that is similar between the two molecules, and (2) may further comprise a sequence that is divergent between the two polynucleotides or amino acid sequences, sequence comparisons between two (or more) molecules are typically performed by comparing sequences of the two molecules over a "comparison window" to identify and compare local regions of sequence similarity. A "comparison window", as used herein, refers to a conceptual segment of at least 18 contiguous nucleotide positions or 6 amino acids wherein a polynucleotide sequence or amino acid sequence may be compared to a reference sequence of at least 18 contiguous nucleotides or 6 amino acid sequences and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions, deletions, substitutions, and the like (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by the local homology algorithm of Smith and Waterman *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman *Proc. Natl. Acad. Sci. (U.S.A.)* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, (Genetics Computer Group, 575 Science Dr., Madison, Wis.), Geneworks, or MacVector software packages), or by

- 33 -

inspection, and the best alignment (i.e., resulting in the highest percentage of homology over the comparison window) generated by the various methods is selected.

- 5 The term "sequence identity" means that two polynucleotide or amino acid sequences are identical (i.e., on a nucleotide-by-nucleotide or residue-by-residue basis) over the comparison window. The term "percentage of sequence identity" is calculated by
- 10 comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, U, or I) or residue occurs in both sequences to yield the number of matched positions,
- 15 dividing the number of matched positions by the total number of positions in the comparison window (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The terms "substantial identity" as used herein denotes a
- 20 characteristic of a polynucleotide or amino acid sequence, wherein the polynucleotide or amino acid comprises a sequence that has at least 85 percent sequence identity, preferably at least 90 to 95 percent sequence identity, more usually at least 99 percent
- 25 sequence identity as compared to a reference sequence over a comparison window of at least 18 nucleotide (6 amino acid) positions, frequently over a window of at least 24-48 nucleotide (8-16 amino acid) positions, wherein the percentage of sequence identity is
- 30 calculated by comparing the reference sequence to the sequence which may include deletions or additions which total 20 percent or less of the reference sequence over

- 34 -

the comparison window. The reference sequence may be a subset of a larger sequence.

As used herein, the twenty conventional amino acids and their abbreviations follow conventional usage. See Immunology - A Synthesis (2nd Edition, E.S. Golub and D.R. Gren, Eds., Sinauer Associates, Sunderland, Mass. (1991)), which is incorporated herein by reference. Stereoisomers (e.g., D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as α -, α -disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional amino acids may also be suitable components for polypeptides of the present invention. Examples of unconventional amino acids include: 4-hydroxyproline, γ -carboxyglutamate, ϵ -N,N,N-trimethyllysine, ϵ -N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, σ -N-methylarginine, and other similar amino acids and imino acids (e.g., 4-hydroxyproline). In the polypeptide notation used herein, the lefthand direction is the amino terminal direction and the righthand direction is the carboxy-terminal direction, in accordance with standard usage and convention.

Similarly, unless specified otherwise, the lefthand end of single-stranded polynucleotide sequences is the 5' end; the lefthand direction of double-stranded polynucleotide sequences is referred to as the 5' direction. The direction of 5' to 3' addition of nascent RNA transcripts is referred to as the transcription direction; sequence regions on the

- 35 -

DNA strand having the same sequence as the RNA and which are 5' to the 5' end of the RNA transcript are referred to as "upstream sequences"; sequence regions on the DNA strand having the same sequence as the RNA and which are 3' to the 3' end of the RNA transcript are referred to as "downstream sequences".

As applied to polypeptides, the term "substantial identity" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 80 percent sequence identity, preferably at least 90 percent sequence identity, more preferably at least 95 percent sequence identity, and most preferably at least 99 percent sequence identity. Preferably, residue positions which are not identical differ by conservative amino acid substitutions. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine,

- 36 -

lysine-arginine, alanine-valine, glutamic-aspartic, and asparagine-glutamine.

As discussed herein, minor variations in the amino acid sequences of antibodies or immunoglobulin molecules are contemplated as being encompassed by the present invention, providing that the variations in the amino acid sequence maintain at least 75%, more preferably at least 80%, 90%, 95%, and most preferably 99%. In particular, conservative amino acid replacements are contemplated. Conservative replacements are those that take place within a family of amino acids that are related in their side chains. Genetically encoded amino acids are generally divided into families: (1) acidic=aspartate, glutamate; (2) basic=lysine, arginine, histidine; (3) non-polar=alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan; and (4) uncharged polar=glycine, asparagine, glutamine, cysteine, serine, threonine, tyrosine. More preferred families are: serine and threonine are aliphatic-hydroxy family; asparagine and glutamine are an amide-containing family; alanine, valine, leucine and isoleucine are an aliphatic family; and phenylalanine, tryptophan, and tyrosine are an aromatic family. For example, it is reasonable to expect that an isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid will not have a major effect on the binding or properties of the resulting molecule, especially if the replacement does not

-37-

involve an amino acid within a framework site. Whether an amino acid change results in a functional peptide can readily be determined by assaying the specific activity of the polypeptide derivative. Assays are
5 described in detail herein. Fragments or analogs of antibodies or immunoglobulin molecules can be readily prepared by those of ordinary skill in the art. Preferred amino- and carboxy-termini of fragments or analogs occur near boundaries of functional domains.
10 Structural and functional domains can be identified by comparison of the nucleotide and/or amino acid sequence data to public or proprietary sequence databases. Preferably, computerized comparison methods are used to identify sequence motifs or predicted protein
15 conformation domains that occur in other proteins of known structure and/or function. Methods to identify protein sequences that fold into a known three-dimensional structure are known. Bowie et al. *Science* 253:164 (1991). Thus, the foregoing examples
20 demonstrate that those of skill in the art can recognize sequence motifs and structural conformations that may be used to define structural and functional domains in accordance with the invention.

25 Preferred amino acid substitutions are those which: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter binding affinities, and (5) confer or modify other
30 physicochemical or functional properties of such analogs. Analogs can include various muteins of a sequence other than the naturally-occurring peptide

- 38 -

sequence. For example, single or multiple amino acid substitutions (preferably conservative amino acid substitutions) may be made in the naturally-occurring sequence (preferably in the portion of the polypeptide outside the domain(s) forming intermolecular contacts. A conservative amino acid substitution should not substantially change the structural characteristics of the parent sequence (e.g., a replacement amino acid should not tend to break a helix that occurs in the parent sequence, or disrupt other types of secondary structure that characterizes the parent sequence). Examples of art-recognized polypeptide secondary and tertiary structures are described in Proteins, Structures and Molecular Principles (Creighton, Ed., W. H. Freeman and Company, New York (1984)); Introduction to Protein Structure (C. Branden and J. Tooze, eds., Garland Publishing, New York, N.Y. (1991)); and Thornton et al. *Nature* 354:105 (1991), which are each incorporated herein by reference.

20

The term "polypeptide fragment" as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion, but where the remaining amino acid sequence is identical to the corresponding positions in the naturally-occurring sequence deduced, for example, from a full-length cDNA sequence. Fragments typically are at least 5, 6, 8 or 10 amino acids long, preferably at least 14 amino acids long, more preferably at least 20 amino acids long, usually at least 50 amino acids long, and even more preferably at least 70 amino acids long. The term "analog" as used herein refers to polypeptides which

30

- 39 -

are comprised of a segment of at least 25 amino acids that has substantial identity to a portion of a deduced amino acid sequence and which has at least one of the following properties: (1) specific binding to a EGF-r, under suitable binding conditions, (2) ability to EGF binding to its receptor, or (3) ability to inhibit EGF-r expressing cell growth in vitro or in vivo.

Typically, polypeptide analogs comprise a conservative amino acid substitution (or addition or deletion) with respect to the naturally-occurring sequence. Analogs typically are at least 20 amino acids long, preferably at least 50 amino acids long or longer, and can often be as long as a full-length naturally-occurring polypeptide.

15

Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed "peptide mimetics" or "peptidomimetics". Fauchere, *J. Adv. Drug Res.* 15:29 (1986); Veber and Freidinger *TINS* p.392 (1985); and Evans et al. *J. Med. Chem.* 30:1229 (1987), which are incorporated herein by reference. Such compounds are often developed with the aid of computerized molecular modeling. Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce an equivalent therapeutic or prophylactic effect. Generally, peptidomimetics are structurally similar to a paradigm polypeptide (i.e., a polypeptide that has a biochemical property or pharmacological activity), such as human antibody, but have one or more peptide linkages

30

-40-

optionally replaced by a linkage selected from the group consisting of: --CH₂NH--, --CH₂S--, --CH₂-CH₂--, --CH=CH-- (cis and trans), --COCH₂--, --CH(OH)CH₂--, and -CH₂SO--, by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) may be used to generate more stable peptides. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus sequence variation may be generated by methods known in the art (Rizo and Gierasch *Ann. Rev. Biochem.* 61:387 (1992), incorporated herein by reference); for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

"Antibody" or "antibody peptide(s)" refer to an intact antibody, or a binding fragment thereof that competes with the intact antibody for specific binding. Binding fragments are produced by recombinant DNA techniques, or by enzymatic or chemical cleavage of intact antibodies. Binding fragments include Fab, Fab', F(ab')₂, Fv, and single-chain antibodies. An antibody other than a "bispecific" or "bifunctional" antibody is understood to have each of its binding sites identical. An antibody substantially inhibits adhesion of a receptor to a counterreceptor when an excess of antibody reduces the quantity of receptor bound to counterreceptor by at least about 20%, 40%, 60% or 80%, and more usually greater than about 85% (as measured in an in vitro competitive binding assay).

-41-

The term "epitope" includes any protein determinant capable of specific binding to an immunoglobulin or T-cell receptor. Epitopic determinants usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics. An antibody is said to specifically bind an antigen when the dissociation constant is $\leq 1 \mu\text{M}$, preferably $\leq 100 \text{ nM}$ and most preferably $\leq 10 \text{ nM}$.

The term "agent" is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials.

As used herein, the terms "label" or "labeled" refers to incorporation of a detectable marker, e.g., by incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotinyl moieties that can be detected by marked avidin (e.g., streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or colorimetric methods). In certain situations, the label or marker can also be therapeutic. Various methods of labeling polypeptides and glycoproteins are known in the art and may be used. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I), fluorescent labels (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic

labels (e.g., horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase), chemiluminescent, biotinyl groups, predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance.

10 The term "pharmaceutical agent or drug" as used herein refers to a chemical compound or composition capable of inducing a desired therapeutic effect when properly administered to a patient. Other chemistry terms herein are used according to conventional usage
15 in the art, as exemplified by The McGraw-Hill Dictionary of Chemical Terms (Parker, S., Ed., McGraw-Hill, San Francisco (1985)), incorporated herein by reference).

20 The term "antineoplastic agent" is used herein to refer to agents that have the functional property of inhibiting a development or progression of a neoplasm in a human, particularly a malignant (cancerous) lesion, such as a carcinoma, sarcoma, lymphoma, or
25 leukemia. Inhibition of metastasis is frequently a property of antineoplastic agents.

As used herein, "substantially pure" means an object species is the predominant species present
30 (i.e., on a molar basis it is more abundant than any other individual species in the composition), and preferably a substantially purified fraction is a

-43-

composition wherein the object species comprises at least about 50 percent (on a molar basis) of all macromolecular species present. Generally, a substantially pure composition will comprise more than 5 about 80 percent of all macromolecular species present in the composition, more preferably more than about 85%, 90%, 95%, and 99%. Most preferably, the object species is purified to essential homogeneity (contaminant species cannot be detected in the 10 composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species.

The term patient includes human and veterinary 15 subjects.

Antibody Structure

The basic antibody structural unit is known to comprise a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having 20 one "light" (about 25 kDa) and one "heavy" chain (about 50-70 kDa). The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal portion of each 25 chain defines a constant region primarily responsible for effector function. Human light chains are classified as kappa and lambda light chains. Heavy chains are classified as mu, delta, gamma, alpha, or epsilon, and define the antibody's isotype as IgM, IgD, 30 IgA, and IgE, respectively. Within light and heavy chains, the variable and constant regions are joined by

-44-

a "J" region of about 12 or more amino acids, with the heavy chain also including a "D" region of about 10 more amino acids. See generally, Fundamental Immunology Ch. 7 (Paul, W., ed., 2nd ed. Raven Press, N.Y. (1989)) (incorporated by reference in its entirety for all purposes). The variable regions of each light/heavy chain pair form the antibody binding site.

Thus, an intact antibody has two binding sites. Except in bifunctional or bispecific antibodies, the two binding sites are the same.

The chains all exhibit the same general structure of relatively conserved framework regions (FR) joined by three hyper variable regions, also called complementarity determining regions or CDRs. The CDRs from the two chains of each pair are aligned by the framework regions, enabling binding to a specific epitope. From N-terminal to C-terminal, both light and heavy chains comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is in accordance with the definitions of Kabat Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md. (1987 and 1991)), or Chothia & Lesk *J. Mol. Biol.* 196:901-917 (1987); Chothia et al. *Nature* 342:878-883 (1989).

A bispecific or bifunctional antibody is an artificial hybrid antibody having two different heavy/light chain pairs and two different binding sites. Bispecific antibodies can be produced by a variety of methods including fusion of hybridomas or linking of Fab' fragments. See, e.g., Songsivilai & Lachmann *Clin. Exp. Immunol.* 79: 315-321 (1990), Kostelny et al. *J. Immunol.* 148:1547-1553 (1992).

-45-

Production of bispecific antibodies can be a relatively labor intensive process compared with production of conventional antibodies and yields and degree of purity are generally lower for bispecific antibodies.

- 5 Bispecific antibodies do not exist in the form of fragments having a single binding site (e.g., Fab, Fab', and Fv).

Preparation of Antibodies

- 10 Antibodies in accordance with the invention are preferably prepared through the utilization of a transgenic mouse that has a substantial portion of the human antibody producing genome inserted but that is rendered deficient in the production of endogenous,
- 15 murine, antibodies. Such mice, then, are capable of producing human immunoglobulin molecules and antibodies and are deficient in the production of murine immunoglobulin molecules and antibodies. Technologies utilized for achieving the same are disclosed in the
- 20 patents, applications, and references disclosed in the Background, herein. In particular, however, a preferred embodiment of transgenic production of mice and antibodies therefrom is disclosed in U.S. Patent Application Serial No. 08/759,620, filed December 3,
- 25 1996, the disclosure of which is hereby incorporated by reference. See also Mendez et al. *Nature Genetics* 15:146-156 (1997), the disclosure of which is hereby incorporated by reference.

- Through use of such technology, we have produced
- 30 fully human monoclonal antibodies to a variety of antigens. Essentially, we immunize XenoMouse™ lines of mice with an antigen of interest, recover lymphatic

- 46 -

cells (such as B-cells) from the mice that express antibodies, fuse such recovered cells with a myeloid-type cell line to prepare immortal hybridoma cell lines, and such hybridoma cell lines are screened and
5 selected to identify hybridoma cell lines that produce antibodies specific to the antigen of interest. We utilized these techniques in accordance with the present invention for the preparation of antibodies specific to EGF-r. Herein, we describe the production
10 of eight hybridoma cell lines that produce antibodies specific to EGF-r. Further, we provide a characterization of the antibodies produced by such cell lines, including nucleotide and amino acid sequence analyses of the heavy and light chains of such
15 antibodies.

The hybridoma cell lines discussed herein are designated E1.1, E2.4, E2.5, E6.2, E6.4, E2.11, E6.3, and E7.6.3. Each of the antibodies produced by the
aforementioned cell lines are fully human IgG2 heavy
20 chains with human kappa light chains. In general, antibodies in accordance with the invention possess very high affinities, typically possessing Kd's of from about 10^{-9} through about 10^{-11} M, when measured by either solid phase and solution phase.

25 As will be appreciated, antibodies in accordance with the present invention can be expressed in cell lines other than hybridoma cell lines. Sequences encoding particular antibodies can be used for transformation of a suitable mammalian host cell.

30 Transformation can be by any known method for introducing polynucleotides into a host cell, including, for example packaging the polynucleotide in

-47-

a virus (or into a viral vector) and transducing a host cell with the virus (or vector) or by transfection procedures known in the art, as exemplified by U.S. Patent Nos. 4,399,216, 4,912,040, 4,740,461, and
5 4,959,455 (which patents are hereby incorporated herein by reference). The transformation procedure used depends upon the host to be transformed. Methods for introduction of heterologous polynucleotides into mammalian cells are well known in the art and include
10 dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei.

15 Mammalian cell lines available as hosts for expression are well known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC), including but not limited to Chinese hamster ovary (CHO) cells, HeLa
20 cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), and a number of other cell lines. Cell lines of particular preference are selected through determining which cell lines have high expression
25 levels and produce antibodies with constitutive EGF-r binding properties.

Antibodies in accordance with the present invention are potent inhibitors of EGF and TGF- α
30 binding to its receptor, EGF-r. Such results are discussed in Examples 5 and 6 and shown in Figures 35 through 38. Consistent with such results, and as shown

-48-

in Figure 39 and discussed in connection with Example 7, antibodies in accordance with the present invention also inhibit the growth of certain human carcinoma cell lines in vitro. Antibodies in accordance with the present invention also prevent the growth of certain human carcinomas in vivo. Such results are shown in Figures 40 through 42 and discussed in connection with Example 8. In Example 9, we demonstrate that antibodies in accordance with the present invention, at least in combination with an antineoplastic agent, will eradicate an existing tumor in an animal. Moreover, antibody therapy, as a monotherapy (i.e., not in combination with an antineoplastic agent) appears possible in accordance with the antibodies in accordance with the present invention, where it did not appear possible in the prior art, for example through the use of the antibody 225. Such results are discussed in connection with Example 9 and shown in Figures 43-44.

The results demonstrated in accordance with the present invention indicate that antibodies in accordance with the present invention possess certain qualities that may make the present antibodies more efficacious than current therapeutic antibodies against EGF-r, e.g., 225. The 225 antibody in clinical development by Imclone is a chimeric IgG1 antibody with an affinity of 2×10^{-10} M, which, while appearing efficacious in combination therapy with an antineoplastic agent, does not appear very efficacious in monotherapy. In contrast, antibodies in accordance with the invention (and particularly the E2.5 and E7.6.3 antibodies of the invention) have significantly

- 49 -

higher affinities ($E2.5:1.6 \times 10^{-11}$ M; $E7.6.3:5.7 \times 10^{-11}$ M) and appear efficacious in monotherapy in addition to combination therapy with an antineoplastic agent and at lower doses than with the C225 antibody.

5

EXAMPLES

The following examples, including the experiments conducted and results achieved are provided for illustrative purposes only and are not to be construed as limiting upon the present invention.

10

Example 1

Generation of Anti-EGF-r-Antibody Producing Hybridomas

Antibodies of the invention were prepared,
15 selected, and assayed in accordance with the present Example.

Immunization and hybridoma generation: XenoMice (8 to 10 weeks old) were immunized intraperitoneally
20 with 2×10^7 A431 (ATCC CRL-7907) cells resuspended in phosphate buffered saline (PBS). This dose was repeated three times. Four days before fusion, the mice received a final injection of cells in PBS. Spleen and lymph node lymphocytes from immunized mice
25 were fused with the non-secretory myeloma NSO-bcl2 line (Ray and Diamond, 1994) and were subjected to HAT selection as previously described (Galfre and Milstein, 1981). A large panel of hybridomas all secreting EGF-r specific human IgG_{2k} (as detected below) antibodies were
30 recovered. As described in Example 2, certain of the

- 50 -

antibodies selected from the panel were selected for their ability to compete with the 225 antibody.

ELISA assay: ELISA for determination of antigen-specific antibodies in mouse serum and in hybridoma supernatants was carried out as described (Coligan et al., 1994) using affinity-purified EGF-r from A431 cells (Sigma, E-3641) to capture the antibodies. The concentrations of human and mouse immunoglobulins were determined using the following capture antibodies: rabbit anti-human IgG (Southern Biotechnology, 6145-01), goat anti-human Igk (Vector Laboratories, AI-3060), mouse anti-human IgM (CGI/ATCC, HB-57), for human gamma, kappa, and mu Ig, respectively, and goat anti-mouse IgG (Caltag, M 30100), goat anti-mouse Igk (Southern Biotechnology, 1050-01), goat anti-mouse IgM (Southern Biotechnology, 1020-01), and goat anti-mouse λ (Southern Biotechnology, 1060-01) to capture mouse gamma, kappa, mu, and lambda Ig, respectively. The detection antibodies used in ELISA experiments were goat anti-mouse IgG-HRP (Caltag, M-30107), goat anti-mouse Igk-HRP (Caltag, M 33007), mouse anti-human IgG2-HRP (Southern Biotechnology, 9070-05), mouse anti-human IgM-HRP (Southern Biotechnology, 9020-05), and goat anti-human kappa-biotin (Vector, BA-3060). Standards used for quantitation of human and mouse Ig were: human IgG₂k (Calbiochem, 400122), human IgMk (Cappel, 13000), mouse IgGk (Cappel 55939), mouse IgMk (Sigma, M-3795), and mouse IgG₃ λ (Sigma, M-9019).

- 51 -

Determination of affinity constants of fully human

Mabs by BIAcore: Affinity measurement of purified human monoclonal antibodies, Fab fragments, or hybridoma supernatants by plasmon resonance was carried out using the BIAcore 2000 instrument, using general procedures outlined by the manufacturers.

Kinetic analysis of the antibodies was carried out using antigens immobilized onto the sensor surface at a low density. Soluble EGF-r purified from A431 cell membranes (Sigma, E-3641) was generally used at a surface density of 228 RU. The dissociation (k_d) and association (k_a) rates were determined using the software provided by the manufacturer (BIA evaluation 2.1).

15 Determination of affinity constants in solution by

ELISA: In order to determine antibody binding affinity in solution by ELISA, various concentrations of the monoclonal antibodies to EGF-r were incubated with EGF-r at a constant concentration until equilibrium was reached. Thereafter, the concentration of the free EGF-r in the reaction solution was determined by an indirect ELISA. Accordingly, the monoclonal antibodies at concentrations of between 3.0×10^{-11} M through 2.7×10^{-7} M were incubated with EGF-r at a concentration of 4×10^{-10} M in 200 μ l of PBS with 0.5% BSA for 15 hrs at room temperature. After incubation, 70 μ l of each mixture was transferred into the wells of 96-well microtiter plates previously coated with the same monoclonal antibody (100 μ l/well, at 2 μ g/ml in coating buffer) and incubated for 15 min at room temperature.

-52-

After washing with washing buffer, the EGF-r retained on the plate was detected by mouse anti-EGF-r-HRP, which binds to the carbohydrate of the EGF-r protein. The concentration of EGF-r was calculated against its standard and used for the calculation of bound and free antibodies in the original antigen-antibody reaction solution. The binding affinity of each monoclonal antibody to EGF-r was calculated using Scatchard analysis.

- 10 Receptor binding assays: The EGF receptor binding assay was carried out with A431 cells or SW948 cells (0.4×10^6 cells per well) which were incubated with varying concentrations of antibodies in PBS binding buffer for 30 minutes at 4°C. 0.1 nM [125 I]EGF
- 15 (Amersham, IM-196) or [125 I]TGF- α (Amersham) was added to each well, and the plates were incubated for 90 min at 4°C. The plates were washed five times, air-dried and counted in a scintillation counter. Anti-EGF-r mouse antibodies 225 and 528 (Calbiochem) were used as
- 20 controls.

EXAMPLE 2

Co-Selection of Anti-EGF-r-Antibodies with the m225 Antibody

- 25 As discussed above, the antibody 225 has been demonstrated to possess a high affinity for, and effective inhibition of the binding of EGF and TGF- α to EGF-r. Thus, we expected that if we selected human antibodies against EGF-r that are prepared in

- 53 -

accordance with the present invention with the antibody 225 in a competition assay, antibodies to the same or similar epitope to which the 225 antibody binds would be selected.

5 Accordingly, we conducted BIAcore assays in which soluble EGF-r purified from A431 cell membranes (Sigma, E-3641) was pretreated with the antibody 225 and thereafter treated with antibodies of the invention. Where antibodies of the invention did not
10 bind, such antibodies of the invention were screened for binding affinity as described above.

In the following Table, affinity measurements for certain of the antibodies selected in this manner are
15 provided:

Table I

	Solid Phase (by BIAcore)				In Solution By ELISA
Hybridoma a	k_{on} ($M^{-1}S^{-1}$)	K_{off} (S^{-1})	K_D (M)	Surface Density [RU]	KD (M)
E1.1	2.3×10^6	1.7×10^{-4}	7.6×10^{-11}	228	1.1×10^{-10}
E2.4	2.8×10^6	9.78×10^{-5}	3.5×10^{-11}	818	1.1×10^{-10}
E2.5	1.2×10^6	1.9×10^{-5}	1.6×10^{-11}	228	3.6×10^{-10}
E2.11	1.9×10^6	3.0×10^{-4}	1.6×10^{-10}	228	1.1×10^{-10}
E7.6.3	2.0×10^6	1.1×10^{-4}	5.7×10^{-11}	228	ND

As will be observed, antibodies selected in this manner possess exceptionally high affinities and binding constants.

5 EXAMPLE 3

Structures of Anti-EGF-r-Antibodies Prepared in Accordance with the Invention

In the following discussion, structural
10 information related to antibodies prepared in accordance with the invention is provided.

In order to analyze structures of antibodies produced in accordance with the invention, we cloned genes encoding the heavy and light chain fragments out
15 of the particular hybridoma. Gene cloning and

-55-

sequencing was accomplished as follows:

Poly(A)⁺ mRNA was isolated from approximately 2 X 10⁵ hybridoma cells derived from immunized XenoMice using a Fast-Track kit (Invitrogen). The generation of random primed cDNA was followed by PCR. Human V_H or human V_κ family specific variable region primers (Marks et. al., 1991) or a universal human V_H primer, MG-30 (CAGGTGCAGCTGGAGCAGTCIGG) (SEQ ID NO:1) was used in conjunction with primers specific for the human Cγ2 constant region (MG-40d; 5'-GCTGAGGGAGTAGAGTCCTGAGGA-3') (SEQ ID NO:2) or Cκ constant region (hκP2; as previously described in Green et al., 1994). Sequences of human Mabs-derived heavy and kappa chain transcripts from hybridomas were obtained by direct sequencing of PCR products generated from poly(A⁺) RNA using the primers described above. PCR products were also cloned into pCRII using a TA cloning kit (Invitrogen) and both strands were sequenced using Prism dye-terminator sequencing kits and an ABI 377 sequencing machine. All sequences were analyzed by alignments to the "V BASE sequence directory" (Tomlinson et al., MRC Centre for Protein Engineering, Cambridge, UK) using MacVector and Geneworks software programs.

25 Hybridoma E1.1

The antibody secreted by the hybridoma E1.1 comprises a human IgG2 antibody having a human kappa light chain. The antibodies were analyzed for structural information related to their heavy chain and light chain gene utilization, as well as their amino acid sequences. Thus, heavy chain V_H, D, and J_H and light chain V_κ and J_κ gene utilization was analyzed and

- 56 -

differences between the coded product and the particular gene utilization was also analyzed. Accordingly, the antibody secreted by the hybridoma E1.1 evidenced the following gene utilization:

5

V_H - 4-31

D - 2

J_H - 5V_K - 018

10

J_K - 4

As reported in the V BASE sequence directory, the amino acid sequence encoded by the V_H 4-31 gene was determined to be:

15

VSGGSISSGGYYWSWIRQHPGKGLEWIGYIYYSGSTYYNPSLKSRVTISVDTSKNQFSLKLSVTAADTAVYYCAR
(SEQ ID NO:19)

As reported in the V BASE sequence directory, the amino acid sequence encoded by the V_K (018) gene was determined to be:

20

TITQASQDISNYLNWYQKPGKAPKLLIYDASNLETGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQYDNLP
(SEQ ID NO:20)

25

Amino acid and nucleotide sequence information respecting the heavy and light chains are provided below in connection with Figures 1-4. Figure 1 is an amino acid sequence of a heavy chain immunoglobulin molecule that is secreted by the hybridoma E1.1.

30

Differences between the sequence encoded by heavy chain variable gene 4-31 and the sequence of the E1.1

- 57 -

secreted heavy chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

5 Figure 2 is a nucleotide sequence of the cDNA encoding the heavy chain immunoglobulin molecule of Figure 1 that was cloned out of the hybridoma E1.1.

Figure 3 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E1.1. Differences between the sequence encoded by light chain variable gene 018 and the sequence of the E1.1 secreted light chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 4 is a nucleotide sequence of the cDNA encoding the kappa light chain immunoglobulin molecule of Figure 3 that was cloned out of the hybridoma E1.1.

20 **Hybridoma E2.4**

The antibody secreted by the hybridoma E2.4 comprises a human IgG2 antibody having a human kappa light chain. The antibodies were analyzed for structural information related to their heavy chain and light chain gene utilization, as well as their amino acid sequences. Thus, heavy chain V_H, D, and J_H and light chain V_K and J_K gene utilization was analyzed and differences between the coded product and the particular gene utilization was also analyzed.

30 Accordingly, the antibody secreted by the hybridoma E2.4 evidenced the following gene utilization:

- 58 -

V_H - 4-31
D - A1/A4
J_H - 3
5 V_K - 018
J_K - 4

Amino acid and nucleotide sequence information
respecting the heavy and light chains are provided
10 below in connection with Figures 5-8. Figure 5 is an
amino acid sequence of a heavy chain immunoglobulin
molecule that is secreted by the hybridoma E2.4.
Differences between the sequence encoded by heavy chain
variable gene 4-31 and the sequence of the E2.4
15 secreted heavy chain are indicated in bold and enlarged
font. The contiguous sequence from CDR1 through CDR3
is indicated by underlining and CDR1, CDR2, and CDR3
sequences are each indicated by double underlining.

Figure 6 is a nucleotide sequence of the cDNA
20 encoding the heavy chain immunoglobulin molecule of
Figure 5 that was cloned out of the hybridoma E2.4.

Figure 7 is an amino acid sequence of a kappa
light chain immunoglobulin molecule that is secreted by
the hybridoma E2.4. Differences between the sequence
25 encoded by light chain variable gene 018 and the
sequence of the E2.4 secreted light chain are indicated
in bold and enlarged font. The contiguous sequence
from CDR1 through CDR3 is indicated by underlining and
CDR1, CDR2, and CDR3 sequences are each indicated by
30 double underlining.

Figure 8 is a nucleotide sequence of the cDNA
encoding the kappa light chain immunoglobulin molecule

- 59 -

of Figure 7 that was cloned out of the hybridoma E2.4.

Hybridoma E2.5

The antibody secreted by the hybridoma E2.5 comprises a human IgG2 antibody having a human kappa
5 light chain. The antibodies were analyzed for structural information related to their heavy chain and light chain gene utilization, as well as their amino acid sequences. Thus, heavy chain V_H , D, and J_H and light chain V_K and J_K gene utilization was analyzed and
10 differences between the coded product and the particular gene utilization was also analyzed. Accordingly, the antibody secreted by the hybridoma E2.5 evidenced the following gene utilization:

15 V_H - 4-31
D - XP1/21-10
 J_H - 4
V K - 018
J K - 2

20

Amino acid and nucleotide sequence information respecting the heavy and light chains are provided below in connection with Figures 9-12. Figure 9 is an amino acid sequence of a heavy chain immunoglobulin
25 molecule that is secreted by the hybridoma E2.5. Differences between the sequence encoded by heavy chain variable gene 4-31 and the sequence of the E2.5 secreted heavy chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3
30 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

- 60 -

Figure 10 is a nucleotide sequence of the cDNA encoding the heavy chain immunoglobulin molecule of Figure 9 that was cloned out of the hybridoma E2.5.

Figure 11 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E2.5. Differences between the sequence encoded by light chain variable gene 018 and the sequence of the E2.5 secreted light chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 12 is a nucleotide sequence of the cDNA encoding the kappa light chain immunoglobulin molecule of Figure 11 that was cloned out of the hybridoma E2.5.

Hybridoma E6.2

The antibody secreted by the hybridoma E6.2 comprises a human IgG2 antibody having a human kappa light chain. The antibodies were analyzed for structural information related to their heavy chain and light chain gene utilization, as well as their amino acid sequences. Thus, heavy chain V_H , D, and J_H and light chain V_k and J_k gene utilization was analyzed and differences between the coded product and the particular gene utilization was also analyzed. Accordingly, the antibody secreted by the hybridoma E6.2 evidenced the following gene utilization:

V_H - 4-31
D - ? (CNTCCCTT)
 J_H - 6

-61-

Vk - 018

Jk - 1

Amino acid and nucleotide sequence information
5 respecting the heavy and light chains are provided
below in connection with Figures 13-16. Figure 13 is
an amino acid sequence of a heavy chain immunoglobulin
molecule that is secreted by the hybridoma E6.2.
Differences between the sequence encoded by heavy chain
10 variable gene 4-31 and the sequence of the E6.2
secreted heavy chain are indicated in bold and enlarged
font. The contiguous sequence from CDR1 through CDR3
is indicated by underlining and CDR1, CDR2, and CDR3
sequences are each indicated by double underlining.

15 Figure 14 is a nucleotide sequence of the cDNA
encoding the heavy chain immunoglobulin molecule of
Figure 13 that was cloned out of the hybridoma E6.2.

Figure 15 is an amino acid sequence of a kappa
light chain immunoglobulin molecule that is secreted by
20 the hybridoma E6.2. Differences between the sequence
encoded by light chain variable gene 018 and the
sequence of the E6.2 secreted light chain are indicated
in bold and enlarged font. The contiguous sequence
from CDR1 through CDR3 is indicated by underlining and
25 CDR1, CDR2, and CDR3 sequences are each indicated by
double underlining.

Figure 16 is a nucleotide sequence of the cDNA
encoding the kappa light chain immunoglobulin molecule
of Figure 15 that was cloned out of the hybridoma E6.2.

30 Hybridoma E6.4

The antibody secreted by the hybridoma E6.4

- 62 -

comprises a human IgG2 antibody having a human kappa light chain. The antibodies were analyzed for structural information related to their heavy chain and light chain gene utilization, as well as their amino acid sequences. Thus, heavy chain V_H, D, and J_H and light chain V_K and J_K gene utilization was analyzed and differences between the coded product and the particular gene utilization was also analyzed. Accordingly, the antibody secreted by the hybridoma E6.4 evidenced the following gene utilization:

V_H - 4-31
D - A1/A4
J_H - 4
V_K - 012
J_K - 2

As reported in the V BASE sequence directory, the amino acid sequence encoded by the V_K 012 gene was determined to be:

TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLT1SSLQPEDFATYYCQQSYSTP
(SEQ ID NO:21)

Amino acid and nucleotide sequence information respecting the heavy and light chains are provided below in connection with Figures 17-20. Figure 17 is an amino acid sequence of a heavy chain immunoglobulin molecule that is secreted by the hybridoma E6.4. Differences between the sequence encoded by heavy chain variable gene 4-31 and the sequence of the E6.4 secreted heavy chain are indicated in bold and enlarged

-63-

font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 18 is a nucleotide sequence of the cDNA
5 encoding the heavy chain immunoglobulin molecule of Figure 17 that was cloned out of the hybridoma E6.4.

Figure 19 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E6.4. Differences between the sequence
10 encoded by light chain variable gene 012 and the sequence of the E6.4 secreted light chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by
15 double underlining.

Figure 20 is a nucleotide sequence of the cDNA encoding the kappa light chain immunoglobulin molecule of Figure 19 that was cloned out of the hybridoma E6.4.

Hybridoma E2.11

20 The antibody secreted by the hybridoma E2.11 comprises a human IgG2 antibody having a human kappa light chain. The antibodies were analyzed for structural information related to their heavy chain and light chain gene utilization, as well as their amino
25 acid sequences. Thus, heavy chain V_H , D, and J_H and light chain V_k and J_k gene utilization was analyzed and differences between the coded product and the particular gene utilization was also analyzed. Accordingly, the antibody secreted by the hybridoma
30 E2.11 evidenced the following gene utilization:

- 64 -

V_H - 4-61
D - XP1/21-10
J_H - 4
V_K - 018
5 J_K - 4

As reported in the V BASE sequence directory, the amino acid sequence encoded by the V_H 4-61 gene was determined to be:

10

VSGGSVSSGSYYWSWIRQPPGKGLEWIGYIYYSGSTNYPNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCAR
(SEQ ID NO:22)

Amino acid and nucleotide sequence information respecting the heavy and light chains are provided below in connection with Figures 21-24. Figure 21 is an amino acid sequence of a heavy chain immunoglobulin molecule that is secreted by the hybridoma E2.11. Differences between the sequence encoded by heavy chain variable gene 4-61 and the sequence of the E2.11 secreted heavy chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 22 is a nucleotide sequence of the cDNA encoding the heavy chain immunoglobulin molecule of Figure 21 that was cloned out of the hybridoma E2.11.

Figure 23 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E2.11. Differences between the sequence encoded by light chain variable gene 018 and the sequence of the E2.11 secreted light chain are

-65-

indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

- 5 Figure 24 is a nucleotide sequence of the cDNA encoding the kappa light chain immunoglobulin molecule of Figure 23 that was cloned out of the hybridoma E2.11.

Hybridoma E6.3

- 10 The antibody secreted by the hybridoma E6.3 comprises a human IgG2 antibody having a human kappa light chain. The antibodies were analyzed for structural information related to their heavy chain and light chain gene utilization, as well as their amino
15 acid sequences. Thus, heavy chain V_H, D, and J_H and light chain V_K and J_K gene utilization was analyzed and differences between the coded product and the particular gene utilization was also analyzed. Accordingly, the antibody secreted by the hybridoma
20 E6.3 evidenced the following gene utilization:

V_H - 4-61

D - 1-2rc

J_H - 4

- 25 V_K - 018

J_K - 4

- Amino acid and nucleotide sequence information respecting the heavy and light chains are provided
30 below in connection with Figures 25-28. Figure 25 is an amino acid sequence of a heavy chain immunoglobulin

-66-

molecule that is secreted by the hybridoma E6.3. Differences between the sequence encoded by heavy chain variable gene 4-61 and the sequence of the E6.3 secreted heavy chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 26 is a nucleotide sequence of the cDNA encoding the heavy chain immunoglobulin molecule of Figure 25 that was cloned out of the hybridoma E6.3.

Figure 27 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E6.3. Differences between the sequence encoded by light chain variable gene 018 and the sequence of the E6.3 secreted light chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 28 is a nucleotide sequence of the cDNA encoding the kappa light chain immunoglobulin molecule of Figure 27 that was cloned out of the hybridoma E6.3.

Hybridoma E7.6.3

The antibody secreted by the hybridoma E7.6.3 comprises a human IgG2 antibody having a human kappa light chain. The antibodies were analyzed for structural information related to their heavy chain and light chain gene utilization, as well as their amino acid sequences. Thus, heavy chain V_H , D, and J_H and light chain V_K and J_K gene utilization was analyzed and differences between the coded product and the

-67-

particular gene utilization was also analyzed. Accordingly, the antibody secreted by the hybridoma E7.6.3 evidenced the following gene utilization:

5 V_H - 4-61
 D - XP4rc-XP1
 J_H - 3
 V_K - 018
 J_K - 4

10

Amino acid and nucleotide sequence information respecting the heavy and light chains are provided below in connection with Figures 29-32. Figure 29 is an amino acid sequence of a heavy chain immunoglobulin molecule that is secreted by the hybridoma E7.6.3. Differences between the sequence encoded by heavy chain variable gene 4-61 and the sequence of the E7.6.3 secreted heavy chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each indicated by double underlining.

Figure 30 is a nucleotide sequence of the cDNA encoding the heavy chain immunoglobulin molecule of Figure 29 that was cloned out of the hybridoma E7.6.3.

25 Figure 31 is an amino acid sequence of a kappa light chain immunoglobulin molecule that is secreted by the hybridoma E7.6.3. Differences between the sequence encoded by light chain variable gene 018 and the sequence of the E7.6.3 secreted light chain are indicated in bold and enlarged font. The contiguous sequence from CDR1 through CDR3 is indicated by underlining and CDR1, CDR2, and CDR3 sequences are each

- 68 -

indicated by double underlining.

Figure 32 is a nucleotide sequence of the cDNA encoding the kappa light chain immunoglobulin molecule of Figure 31 that was cloned out of the hybridoma

5 E7.6.3.

EXAMPLE 4

Analysis of Heavy and Light Chain Amino Acid Substitutions

10 Figure 33 provides a comparison of specific anti-EGF-r antibody heavy chain amino acid sequence comparisons with the amino acid sequence of the particular V_H gene which encodes the heavy chain of the particular antibody. Figure 34 provides a similar
15 comparison of specific anti-EGF-r antibody light chain amino acid sequence comparisons with the amino acid sequence of the particular V_k gene which encodes the light chain of the particular antibody. As will be observed, there are several remarkably conserved amino
20 acid substitutions amongst the heavy and light chain sequences. In particular, in the heavy chains of the antibodies, all of the heavy chain molecules are encoded by V_H 4 family genes and have a Glycine in position 10 in V_H 4-31 encoded antibodies and Serine in
25 position 10 in V_H 4-61 encoded antibodies are each substituted with an Aspartic Acid. Also in the V_H 4-31 heavy chains, all but one of the antibodies includes a Serine in position 7 substitution to Asparagine. A similar, though not quite as predominant substitution
30 is observed in position 35, where a Serine in two of the V_H 4-31 encoded antibodies and two of the V_H 4-61

- 69 -

encoded antibodies is substituted with an Asparagine. Also, in two of the V_H 4-31 encoded antibodies and two of the V_H 4-61 encoded antibodies there are substitutions at position 28, where in each case, a
5 Tyrosine is substituted with a Serine (E2.4) or a Histidine (E6.4, E2.11, and E7.6.3). Five of the antibodies, three of the V_H 4-31 encoded antibodies and two of the V_H 4-61 encoded antibodies, possess Valine to Leucine (E2.4 and E2.11) or Isoleucine (E2.5, E6.2, and
10 E7.6.3) at position 50.

In connection with the kappa light chains amino acid sequences, all of the sequences are encoded by V_K I family genes, with seven of the molecules being encoded by 018 genes and one (E6.4) being encoded by an
15 012 gene. There is a high degree of homology between the 012 and 018 gene products, as evidenced when the E6.4 molecule is compared with the 018 gene product, along with the other molecules, in Figure 34. The E6.4 molecule possesses only two substitutions relative to
20 the 012 gene product, as shown in Figure 19, and only 13 substitutions relative to the 018 gene product. All of the antibodies possess a substitution at position 74 in CDR3 where an Asparagine is substituted with a Serine (E1.1, E2.5, E2.11, and E6.3), Histidine (E2.4,
25 E6.2, and E7.6.3), or Arginine (E6.4). The remainder of the substitutions are less highly conserved. However, a number of the antibodies appear to possess substitutions within the CDR's. However, it is interesting to note that E7.6.3, which is an antibody
30 with very high affinities, possesses no amino acid substitutions in the light chain amino acid sequence until just proximal to CDR3 and within CDR3.

- 70 -

It will be appreciated that each of the above-identified amino acid substitutions exist in close proximity to or within a CDR. Such substitutions would appear to bear some effect upon the binding of the antibody to the EGF receptor molecule. Further, such substitutions could have significant effect upon the affinity of the antibodies.

As was discussed above, anti-EGF-r antibodies have been demonstrated to possess certain anti-tumor activities. The following experiments were carried out in order to determine if antibodies in accordance with the present invention possessed such anti-tumor activities.

EXAMPLE 5

Blockage of EGF and TGF- α Binding to Human Epidermoid Carcinoma A431 Cells by Human Anti-EGF-r Antibodies in vitro

An in vitro assay was conducted to determine if antibodies in accordance with the present invention were capable of blocking EGF binding to a human carcinoma cell line. The experiment was conducted to compare the binding of antibodies in accordance with the invention with the murine monoclonal antibody 225 which, as discussed above, has previously demonstrated anti-cancer activity.

In this example, the human epidermoid carcinoma A431 cell line was utilized. The A431 cell line is known for its high expression level of EGF-r (about 2 X

-71-

10⁶ EGF-r molecules per cell). Therefore, higher concentrations of anti-EGF-r antibodies are required to saturate all of the binding sites. The results from this example are shown in Figure 35. In the Figure, blockage of I¹²⁵ labeled EGF binding to human epidermoid carcinoma A431 cells by a human anti-EGF-r antibody in vitro is demonstrated. In the Figure, (□) depicts the results achieved by the anti-EGF-r antibody in accordance with the invention (E7.6.3), (O) depicts the results achieved by the murine monoclonal antibody 225, and (s) depicts the results achieved by a control, nonspecific, human IgG2 antibody.

Figure 36 shows inhibition of EGF binding to human epidermoid carcinoma A431 cells by a panel of human anti-EGF-r antibodies in accordance with the invention in vitro when compared to the 225, 528, and nonspecific human IgG2 controls. In the Figure, (□) depicts the results achieved by the murine monoclonal antibody 225, (O) depicts the results achieved by the murine monoclonal antibody 528, (t) depicts the results achieved using the E1.1 antibody in accordance with the invention, (s) depicts the results achieved using the E2.4 antibody in accordance with the invention, (4) depicts the results achieved using the E2.5 antibody in accordance with the invention, (3) depicts the results achieved using the E2.6 antibody in accordance with the invention, (u) depicts the results achieved using the E2.11 antibody in accordance with the invention, and (—) depicts the results achieved using a control, nonspecific human IgG2 antibody.

The results indicate that antibodies in accordance with the invention may block EGF binding to surface

- 72 -

expressed EGF-r on A431 cells better than the 225 and 528 antibodies. Antibodies in accordance with the invention appear to begin inhibiting binding at an 8 nM concentration as compared to a 10 nM concentration for the 225 antibody.

In connection with inhibition of TGF- α binding, similar efficacy is observed through use of antibodies in accordance with the invention when compared to the 225 antibody. Figure 37 shows inhibition of TGF- α binding to human epidermoid carcinoma A431 cells by human anti-EGF-r antibodies in vitro, where (\square) depicts the results achieved by the murine monoclonal antibody 225, (u) depicts the results achieved using the E6.2 antibody in accordance with the invention, (l) depicts the results achieved using the E6.3 antibody in accordance with the invention, (s) depicts the results achieved using the E7.2 antibody in accordance with the invention, (n) depicts the results achieved using the E7.10 antibody in accordance with the invention, (t) depicts the results achieved using the E7.6.3, and (') depicts the results achieved using a control, nonspecific human IgG2 antibody.

The results indicate that antibodies in accordance with the invention may block TGF- α binding to surface expressed EGF-r on A431 cells better than the 225 antibody. Antibodies in accordance with the invention appear to begin inhibiting binding at an 0.1 nM concentration as compared to a 1 nM concentration for the 225 antibody.

30

EXAMPLE 6

Blockage of EGF Binding to Human Colon Adenocarcinoma

SW948 Cells by Human Anti-EGF-r Antibodies in vitro

Another in vitro assay was conducted to determine if antibodies in accordance with the present invention were capable of blocking EGF binding to yet another human carcinoma cell line. The experiment was conducted to compare the binding of antibodies in accordance with the invention with the murine monoclonal antibody 225 which, as discussed above, has previously demonstrated anti-cancer activity.

In this example, the human colon adenocarcinoma SW948 cell line was utilized. In contrast to the A431 cell line, the SW948 cell line has relatively low expression of EGF-r on its surface (about 40,000 molecules per cell). Therefore, less of the anti-EGF-r antibodies are required to saturate all of the binding sites of the receptors on the cells. The results from this example are shown in Figure 38. In the Figure, blockage of I^{125} labeled EGF binding to human colon adenocarcinoma SW948 cells by a human anti-EGF-r antibody in vitro is demonstrated. In the Figure, (m) depicts the results achieved by an anti-EGF-r antibody in accordance with the invention (E7.6.3), (\square) depicts the results achieved by the murine monoclonal antibody 225, and (s) depicts the results achieved by a control, nonspecific, human IgG2 antibody.

The results indicate that the antibody in accordance with the invention blocks EGF binding to SW948 cells at least as well as the 225 antibody. In fact, the curve is slightly improved with respect to the antibody in accordance with the invention, showing inhibition at lower concentrations than the 225

antibody.

EXAMPLE 7

Inhibition of Human Colon Adenocarcinoma SW948 Cell

5 Growth by Human Anti-EGF-r Antibodies in vitro

We also conducted an in vitro assay to determine whether and to what degree, as compared to the 225 antibody, antibodies in accordance with the invention were capable of inhibiting cancer cell growth. The experiment was conducted to compare the inhibition by antibodies in accordance with the invention with the inhibition by the murine monoclonal antibody 225 which, as discussed above, has previously demonstrated anti-cancer activity.

In this example, the human colon adenocarcinoma SW948 cell line was utilized. In our hands, only the SW948 cell line showed EGF-dependent cell growth. In contrast, the A431 cell line showed growth inhibition in the presence of EGF in vitro. The results are shown in Figure 39 where it is demonstrated that human anti-EGF-r antibodies in accordance with the present invention inhibit the growth of SW948 cells in vitro. In the Figure, (m) depicts the results achieved by an anti-EGF-r antibody in accordance with the invention (E7.6.3), (□) depicts the results achieved by the murine monoclonal antibody 225, and (s) depicts the results achieved by a control, nonspecific, human IgG2 antibody.

30 The results indicate that the antibody in accordance with the invention inhibits growth of SW948 cells at least as well as the 225 antibody. In fact,

- 75 -

the curve is slightly improved with respect to the antibody in accordance with the invention, showing an apparent 100% inhibition in cell growth at approximately 100 µg/ml whereas the 225 antibody appears to plateau at an inhibition level between 80 to 90% in the same dosage range.

EXAMPLE 8

Inhibition of Human Epidermoid Carcinoma Growth in Nude Mice by Human Anti-EGF-r Antibodies in vivo

In the present experiment, we sought to determine if antibodies in accordance with the present invention were capable of inhibiting tumor cell growth in vivo.

In the experiment, nude mice at the age of 8 weeks were inoculated subcutaneously with the human epidermoid carcinoma A431 cell line. Mice were injected with 5×10^6 A431 cells. One of two dosages of an antibody in accordance with the invention or one of two controls was injected intraperitoneally on the same day when the A431 cells were inoculated. Three administrations of either antibody or control followed and mice were followed for subcutaneous tumor formation and size. The dosages of antibody utilized were either 1.0 mg or 0.2 mg. The controls were either phosphate buffered saline or a nonspecific human IgG2 antibody.

The results from this experiment are shown in Figure 40. In the Figure, the inhibition of human epidermoid carcinoma cell growth in nude mice through use of human anti-EGF-r antibodies in accordance with the invention in vivo is evident. In the Figure, (s) depicts the results achieved with a dosage of 1.0 mg of

-76-

a human anti-EGF-r antibody in accordance with the present invention (E7.6.3) (n=5), (t) depicts the results achieved with a dosage of 0.2 mg of the E.7.6.3 antibody (n=4), (□) depicts the results achieved by a control, nonspecific, human IgG2 antibody (n=6), and (m) depicts the results achieved utilizing phosphate buffered saline as a control (n=6).

No tumor growth was observed in animals treated with the E7.6.3 antibody whereas control animals grew significant tumors within 25 days of tumor cell inoculation.

In the same experiment, three antibodies in accordance with the invention were compared. The results are shown in Figure 41. Each of the antibodies in accordance with the present invention, E7.6.3 at 1 mg in 5 mice and 0.2 mg in 4 mice, E2.5 at 1 mg in 3 mice and 0.2 mg in 3 mice, and E1.1 at 1 mg in 3 mice, demonstrated inhibition of the human epidermoid carcinoma formation in the mice in comparison to controls. All of the control animals (including 6 PBS-treated animals and 6 human IgG2-treated animals) developed significant tumors within 19 days of inoculation whereas none of the the animals treated with the human anti-EGF-r antibodies in accordance with the invention developed tumors within 19 days of inoculation.

Figure 42 shows the results of following the animals from this above-mentioned same experiment for 130 days post inoculation with the human epidermoid carcinoma. The results from this experiment are shown in Figure 42. In the Figure, it will be observed that all of the control mice had developed tumors within 20

-77-

days of tumor cell inoculation. In contrast, the first mouse treated with an antibody in accordance with the present invention to develop a tumor was on day 70. By day 130, only 4 out of 15 of the experimental animals had developed tumors. Interestingly, none of the experimental animals treated with the 0.2 mg dosage of the E2.5 antibody developed tumors within the test period.

The above experiment in connection with this Example 8 demonstrate that antibodies in accordance with the present invention if administered contemporaneously with the inoculation of a tumor cell line appear to almost entirely prevent the initiation of tumor cell growth and initiation of the tumor. Moreover, it will be observed that the inhibitory effect on tumor cell growth appears long-lasting.

EXAMPLE 9

Eradication of Human Epidermoid Carcinoma Growth in Nude Mice by Human Anti-EGF-r Antibodies in vivo

While preventing tumor cell growth and/or establishment of a tumor, as discussed above in connection with the preceding example, is a positive finding, from a therapeutic point of view, eradication of an established tumor is also highly desirable. Accordingly, in the following experiments we examined whether antibodies in accordance with the invention were capable of eradicating an established tumor in a mammal. Previous data generated in connection with the 225 antibody indicated that in order to effectively eradicate an established tumor through use of the 225 antibody it was necessary to complement treatment with

- 78 -

an antineoplastic agent. Thus, in connection with our experiments, we looked at antibody treatment both alone and in combination with antineoplastic agent treatment.

In the experiment, nude mice were inoculated subcutaneously with 5×10^6 A431 human epidermoid carcinoma cells on day 0. Mice were treated with either antibodies, chemotherapeutic agents, and/or controls after the tumor had an opportunity to become established (size ≥ 0.4 cm³). Treatments were begun and continued on days 5, 8, 10, 14, 16, and 21, with chemotherapies being administered only on days 5 and 6. Therapies consisted of an antibody in accordance with the invention (E7.6.3), the antineoplastic agent doxorubicin, and a combination of antibody and doxorubicin. Controls were phosphate buffered saline or a nonspecific human IgG2 antibody. Each treatment group consisted of 5 animals. The data generated from the experiments are shown in Figure 43, where (s) depicts the results achieved with a dosage of 1 mg of a human anti-EGF-r antibody in accordance with the present invention (E7.6.3) (n=5), (5) depicts the results achieved with a dosage of 125 μ g of doxorubicin, (V) depicts the results achieved with a dosage of 1 mg of a human anti-EGF-r antibody in accordance with the present invention (E7.6.3) in combination with a dosage of 125 μ g of doxorubicin, (n) depicts the results achieved by a control, nonspecific, human IgG2 antibody, and (u) depicts the results achieved utilizing phosphate buffered saline as a control.

As will be observed, administration of the E7.6.3 antibody in combination with doxorubicin resulted in

- 79 -

complete eradication tumor growth. Further, tumor growth was completely arrested through administration of the E7.6.3 antibody alone.

In a similar experiment, the results of which are shown in Figure 44, following inoculation with the tumor, five mice were treated with 0.5 mg of the E2.5 antibody on days 5, 8, 10, 14, 16, and 21 and five mice were treated with a combination of the E2.5 antibody administered on days 5, 8, 10, 14, 16, and 21 and doxorubicin administered on days 5 and 6. In the Figure, (u) depicts the results achieved with a dosage of 0.5 mg of a human anti-EGF-r antibody in accordance with the present invention (E2.5), (n) depicts the results achieved with a dosage of 125 µg of doxorubicin, (s) depicts the results achieved with a dosage of 0.5 mg of a human anti-EGF-r antibody in accordance with the present invention (E2.5) in combination with a dosage of 125 µg of doxorubicin, (5) depicts the results achieved utilizing phosphate buffered saline as a control, and (V) depicts the results achieved utilizing a control, nonspecific, human IgG2 antibody.

As will be observed, administration of the E2.5 antibody by itself, or in combination with doxorubicin, resulted in near complete eradication of tumors in the mice.

EXAMPLE 10

Human Clinical Trials for the Treatment and Diagnosis of Human Carcinomas through use of Human Anti-EGF-r Antibodies in vivo

Introduction

Antibodies in accordance with the present invention are indicated in the treatment of certain solid tumors. Based upon a number of factors, including EGF-r expression levels, among others, the following tumor types appear to present preferred indications: breast, ovarian, colon, prostate, bladder and non-small cell lung cancer. In connection with each of these indications, three clinical pathways appear to offer distinct potentials for clinical success:

Adjunctive therapy: In adjunctive therapy, patients would be treated with antibodies in accordance with the present invention in combination with a chemotherapeutic or antineoplastic agent and/or radiation therapy. The primary targets listed above will be treated under protocol by the addition of antibodies of the invention to standard first and second line therapy. Protocol designs will address effectiveness as assessed by reduction in tumor mass as well as the ability to reduce usual doses of standard chemotherapy. These dosage reductions will allow additional and/or prolonged therapy by reducing dose-related toxicity of the chemotherapeutic agent. Prior art anti-EGF-r antibodies have been, or are being, utilized in several adjunctive clinical trials in combination with the chemotherapeutic or antineoplastic agents adriamycin (C225: advanced prostate carcinoma), cisplatin (C225: advanced head and neck and lung carcinomas), taxol (C225: breast cancer), and doxorubicin (C225: preclinical).

Monotherapy: In connection with the use of the

- 81 -

antibodies in accordance with the present invention in monotherapy of tumors, the antibodies will be administered to patients without a chemotherapeutic or antineoplastic agent. Preclinical results generated through use of antibodies in accordance with the present invention and discussed herein have demonstrated similar results with both adjunctive therapy and/or as a stand-alone therapy. Moreover, monotherapy has apparently been conducted clinically in end stage cancer patients with extensive metastatic disease. Patients appeared to show some disease stabilization. Id. Trials will be designed to demonstrate an effect in refractory patients with (cancer) tumor.

15 Imaging Agent: Through binding a radionuclide (e.g., yttrium (^{90}Y)) to antibodies in accordance with the present invention, it is expected that radiolabeled antibodies in accordance with the present invention can be utilized as a diagnostic, imaging agent. In such a role, antibodies of the invention will localize to both solid tumors, as well as, metastatic lesions of cells expressing the EGF receptor. In connection with the use of the antibodies of the invention as imaging agents, the antibodies can be used in assisting surgical treatment of solid tumors, as both a pre-surgical screen as well as a post operative follow to determine what tumor remain and/or returns. An (^{111}In)-C225 antibody has been used as an imaging agent in a Phase I human clinical trial in patients having unresectable squamous cell lung carcinomas. Divgi et al. *J. Natl. Cancer Inst.* 83:97-104 (1991). Patients were followed with standard anterior and posterior

- 82 -

gamma camera. Preliminary data indicated that all primary lesions and large metastatic lesions were identified, while only one-half of small metastatic lesions (under 1 cm) were detected.

5 Dose and Route of Administration

While specific dosing for antibodies in accordance with the invention has not yet been determined, certain dosing considerations can be determined through comparison with the similar product (ImClone C225) that is in the clinic. The C225 antibody is typically being administered with doses in the range of 5 to 400 mg/m², with the lower doses used only in connection with the safety studies. Antibodies in accordance with the invention have a one-log higher affinity than the C225 antibody. Further, antibodies in accordance with the present invention are fully human antibodies, as compared to the chimeric nature of the C225 antibody and, thus, antibody clearance would be expected to be slower. Accordingly, we would expect that dosing in patients with antibodies in accordance with the invention can be lower, perhaps in the range of 50 to 300 mg/m², and still remain efficacious. Dosing in mg/m², as opposed to the conventional measurement of dose in mg/kg, is a measurement based on surface area and is a convenient dosing measurement that is designed to include patients of all sizes from infants to adults.

Three distinct delivery approaches are expected to be useful for delivery of the antibodies in accordance with the invention. Conventional intravenous delivery will presumably be the standard delivery technique for

- 83 -

the majority of tumors. However, in connection with tumors in the peritoneal cavity, such as tumors of the ovaries, biliary duct, other ducts, and the like, intraperitoneal administration may prove favorable for obtaining high dose of antibody at the tumor and to minimize antibody clearance. In a similar manner certain solid tumors possess vasculature that is appropriate for regional perfusion. Regional perfusion will allow the obtention of a high dose of the antibody at the site of a tumor and will minimize short term clearance of the antibody.

Clinical Development Plan (CDP)

Overview: The CDP will follow and develop treatments of anti-EGF-r antibodies in accordance with the invention in connection with adjunctive therapy, monotherapy, and as an imaging agent. Trials will be initially utilized to demonstrate safety and will thereafter be utilized to address efficacy in repeat doses. Trails will be open label comparing standard chemotherapy with standard therapy plus antibodies in accordance with the invention. As will be appreciated, one criteria that can be utilized in connection with enrollment of patients can be EGF-r expression levels of patient tumors as determined in biopsy.

As with any protein or antibody infusion based therapeutic, safety concerns are related primarily to (i) cytokine release syndrome, i.e., hypotension, fever, shaking, chills, (ii) the development of an immunogenic response to the material (i.e., development of human antibodies by the patient to the human antibody therapeutic, or HABA response), and (iii)

- 84 -

toxicity to normal cells that express the EGF receptor, e.g., hepatocytes which express EGF-r. Standard tests and follow up will be utilized to monitor each of these safety concerns. In particular, liver function will be
5 monitored frequently during clinical trials in order to assess damage to the liver, if any.

Human Clinical Trial: Adjunctive Therapy with
Human Anti-EGF-r Antibody and Chemotherapeutic
Agent

10 A phase I human clinical trial will be initiated to assess the safety of six intravenous doses of a human anti-EGF-r antibody in accordance with the invention in connection with the treatment of a solid tumor, e.g., breast cancer. In the
15 study, the safety of single doses of antibodies in accordance with the invention when utilized as an adjunctive therapy to an antineoplastic or chemotherapeutic agent, such as cisplatin, topotecan, doxorubicin, adriamycin, taxol, or the
20 like, will be assessed. The trial design will include delivery of six, single doses of an antibody in accordance with the invention with dosage of antibody escalating from approximately about 25 mg/m² to about 275 mg/m² over the course of
25 the treatment in accordance with the following schedule:

- 85 -

	Day 0	Day 7	Day 14	Day 21	Day 28	Day 35
Mab Dose	25	75	125	175	225	275
	mg/m ²	mg/m ²	mg/m ²	mg/m ²	mg/m ²	mg/m ²
Chemotherapy (standard dose)	+	+	+	+	+	+

Patients will be closely followed for one-week
 5 following each administration of antibody and
 chemotherapy. In particular, patients will be
 assessed for the safety concerns mentioned above: (i)
 cytokine release syndrome, i.e., hypotension, fever,
 shaking, chills, (ii) the development of an
 10 immunogenic response to the material (i.e.,
 development of human antibodies by the patient to the
 human antibody therapeutic, or HAHA response), and
 (iii) toxicity to normal cells that express the EGF
 receptor, e.g., hepatocytes which express EGF-r.
 15 Standard tests and follow up will be utilized to
 monitor each of these safety concerns. In particular,
 liver function will be monitored frequently during
 clinical trials in order to assess damage to the
 liver, if any.

20 Patients will also be assessed for clinical
 outcome, and particularly reduction in tumor mass as
 evidenced by MRI or other imaging.

Assuming demonstration of safety and an
 indication of efficacy, Phase II trials would likely
 25 be initiated to further explore the efficacy and

-86-

determine optimum dosing.

Human Clinical Trial: Monotherapy with Human Anti-EGF-r Antibody

- 5 Assuming that the antibodies in accordance with the present invention appear safe in connection with the above-discussed adjunctive trial, a human clinical trial to assess the efficacy and optimum dosing for monotherapy. Such trial could be accomplished, and
10 would entail the same safety and outcome analyses, to the above-described adjunctive trial with the exception being that patients will not receive chemotherapy concurrently with the receipt of doses of antibodies in accordance with the invention.

15 Human Clinical Trial: Diagnostic Imaging with Anti-EGF-r Antibody

- Once again, assuming that the adjunctive therapy discussed above appears safe within the safety criteria discussed above, a human clinical
20 trial can be conducted concerning the use of antibodies in accordance with the present invention as a diagnostic imaging agent. It is expected that the protocol would be designed in a substantially similar manner to that described in Divgi et al. J.
25 Natl. Cancer Inst. 83:97-104 (1991).

INCORPORATION BY REFERENCE

 All references cited herein, including patents, patent applications, papers, text books, and the

- 87 -

like, and the references cited therein, to the extent that they are not already, are hereby incorporated herein by reference in their entirety. In addition, the following references are also incorporated by
5 reference herein in their entirety, including the references cited in such references:

Abertsen et al., "Construction and characterization of a yeast artificial chromosome library containing seven haploid human genome equivalents." *Proc. Natl.*
10 *Acad. Sci.* 87:4256 (1990).

Anand et al., "Construction of yeast artificial chromosome libraries with large inserts using fractionation by pulsed-field gel electrophoresis."
Nucl. Acids Res. 17:3425-3433 (1989).

15 Berman et al. "Content and organization of the human Ig VH locus: definition of three new V_H families and linkage to the Ig C_H locus." *EMBO J.* 7:727-738 (1988).

Brezinschek et al., "Analysis of the heavy chain
20 repertoire of human peripheral B-cells using single-cell polymerase chain reaction." *J. Immunol.* 155:190-202 (1995).

Brownstein et al., "Isolation of single-copy human genes from a library of yeast artificial chromosome
25 clones." *Science* 244:1348- 1351 (1989).

Bruggeman et al. *PNAS USA* 86:6709-6713 (1989).

Bruggemann et al., "Human antibody production in transgenic mice: expression from 100 kb of the human IgH locus." *Eur. J. Immunol.* 21:1323-1326 (1991).

- 5 Bruggeman, M. and Neuberger, M.S. in *Methods: A companion to Methods in Enzymology* 2:159-165 (Lerner et al. eds. Academic Press (1991)).

Bruggemann, M. and Neuberger, M.S. "Strategies for expressing human antibody repertoires in transgenic
10 mice." *Immunology Today* 17:391-397 (1996).

Chen et al. "Immunoglobulin gene rearrangement in B-cell deficient mice generated by targeted deletion of the J_H locus" *International Immunology* 5:647-656 (1993)

- 15 Choi et al. "Transgenic mice containing a human heavy chain immunoglobulin gene fragment cloned in a yeast artificial chromosome" *Nature Genetics* 4:117-123 (1993)

Coligan et al., Unit 2.1, "Enzyme-linked
20 immunosorbent assays," in *Current protocols in immunology* (1994).

Cook, G.P. and Tomlinson, I.M., "The human immunoglobulin V_H repertoire." *Immunology Today* 16:237-242 (1995).

- Cox et al., "A directory of human germ-line Vx segments reveals a strong bias in their usage." *Eur. J. Immunol.* 24:827-836 (1994).
- Dariavach et al., "The mouse IgH 3'-enhancer." *Eur. J. Immunol.* 21:1499-1504 (1991).
- Den Dunnen et al., "Reconstruction of the 2.4 Mb human DMD-gene by homologous YAC recombination." *Human Molecular Genetics* 1:19-28 (1992).
- Feeney, A.J. "Lack of N regions in fetal and neonatal mouse immunoglobulin V-D-J junctional sequences." *J. Exp. Med.* 172:137-1390 (1990).
- Fishwild et al., "High-avidity human IgGκ monoclonal antibodies from a novel strain of minilocus transgenic mice." *Nature Biotech.* 14:845-851 (1996).
- 15 Flanagan, J.G. and Rabbitts, T.H., "Arrangement of human immunoglobulin heavy chain constant region genes implies evolutionary duplication of a segment containing g, e, and a genes." *Nature* 300:709-713 (1982).
- 20 Galfre, G. and Milstein, C., "Preparation of monoclonal antibodies: strategies and procedures." *Methods Enzymol.* 73:3-46 (1981).
- Gemmell et al., "Protocols for pulsed field gel electrophoresis: Separation and detection of large

-90-

DNA molecules." *Advances in Genome Biology* 1:217-251 (1991).

Gill et al., "Monoclonal anti-epidermal growth factor receptor antibodies which are inhibitors of epidermal growth factor binding and antagonists of epidermal growth factorstimulated tyrosine protein kinase activity." *J. Biol. Chem.* 259:7755 (1984).

Green et al., "Antigen-specific human monoclonal antibodies from mice engineered with human Ig heavy and light chain YACs." *Nature Genetics* 7:13-21 (1994).

Hermanson et al., "Rescue of end fragments of yeast artificial chromosomes by homologous recombination in yeast." *Nucleic Acids Res.* 19:4943-4948 (1991).

15 Huber et al., "The human immunoglobulin κ locus. Characterization of the partially duplicated L regions." *Eur. J. Immunol.* 23:2860-2967 (1993).

Jakobovits, A., "Humanizing the mouse genome." *Current Biology* 4:761-763 (1994).

20 Jakobovits, A., "Production of fully human antibodies by transgenic mice." *Current Opinion in Biotechnology* 6:561-566 (1995).

Jakobovits et al., "Germ-line transmission and expression of a human-derived yeast

- 91 -

artificial-chromosome." Nature 362:255-258 (1993).

Jakobovits, A. et al., "Analysis of homozygous mutant chimeric mice: Deletion of the immunoglobulin heavy-chain joining region blocks B-cell development and antibody production." Proc. Natl. Acad. Sci. USA 90:2551-2555 (1993).

Kawamoto et al., "Growth stimulation of A431 cells by epidermal growth factor: Identification of high affinity receptors for EGF by an anti-receptor monoclonal antibody." Proc. Nat. Acad. Sci., USA 80:1337-1341 (1983).

Lonberg et al., "Antigen-specific human antibodies from mice comprising four distinct genetic modifications." Nature 368:856-859 (1994).

15 Lusti-Marasimhan et al., "Mutation of Leu25 and Val27 introduces CC chemokine activity into interleukin-8." J. Biol. Chem. 270:2716-2721 (1995).

Marks et al., "Oligonucleotide primers for polymerase chain reaction amplification of human immunoglobulin variable genes and design of family-specific oligonucleotide probes." Eur. J. Immunol. 21:985-991 (1991).

Matsuda et al., "Structure and physical map of 64 variable segments in the 3' 0.8-megabase region of the human immunoglobulin heavy-chain locus." Nature 25

Genetics 3:88-94 (1993).

Max, E. Molecular genetics of immunoglobulins.

Fundamental Immunology. 315-382 (Paul, WE, ed., New York: Raven Press (1993)).

- 5 Mendez et al., "A set of YAC targeting vectors for the interconversion of centric and acentric arms." Cold Spring Harbor Laboratory Press, Genome Mapping and Sequencing meeting, 163 (1993).

- Mendez et al., "Analysis of the structural integrity
10 of YACs comprising human immunoglobulin genes in yeast and in embryonic stem cells." Genomics 26:294-307 (1995).

- Ray, S. and Diamond, B., "Generation of a fusion
partner to sample the repertoire of Splenic B-cells
15 destined for apoptosis." Proc. Natl. Acad. Sci. USA 91:5548-5551 (1994).

Sato et al., "Biological effects in vitro of monoclonal antibodies to human epidermal growth factor receptors" Mol. Biol. Med. 1:511-529 (1983).

- 20 Schiestl, R.H. and Gietz, R.D., "High efficiency transformation of intact yeast cells using stranded nucleic acids as a carrier." Curr. Genet. 16:339-346 (1989).

Sherman et al., "Laboratory Course Manual for Methods

- 93 -

in Yeast Genetics." (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1986)).

Silverman et al., "Meiotic recombination between yeast artificial chromosomes yields a single clone
5 containing the entire BCL2 protooncogene." Proc. Natl. Acad. Sci. USA 87:9913-9917 (19__).

Srivastava, A. and Schlessinger, D., "Vectors for inserting selectable markers in vector arms and human DNA inserts of yeast artificial chromosomes (YACs)."
10 Gene 103:53-59 (1991).

Taylor et al., "A transgenic mouse that expresses a diversity of human sequence heavy and light chain immunoglobulins." Nucleic Acids Research 20:6287-6295 (1992).

15 Taylor et al., "Human immunoglobulin transgenes undergo rearrangement, somatic mutation and class switching in mice that lack endogenous IgM." International Immunology 6:579-591 (1994).

Tuailon et al., "Human immunoglobulin heavy-chain
20 minilocus recombination in transgenic mice: gene-segment use in m and g transcripts." Proc. Natl. Acad. Sci. USA 90:3720-3724 (1993).

Tuailon et al. "Analysis of direct and inverted DJ_H rearrangements in a human Ig heavy chain transgenic
25 minilocus" J. Immunol. 154:6453-6465 (1995)

- 94 -

Vaughan et al., "Human antibodies with subnanomolar affinities isolated from a large non-immunized phage display library." Nature Biotech. 14:309-314 (1996).

Wagner et al., "The diversity of antigen-specific
5 monoclonal antibodies from transgenic mice bearing
human immunoglobulin gene miniloci." Eur. J.
Immunol. 24:2672-2681 (1994).

Weichhold et al., "The human immunoglobulin κ locus
consists of two copies that are organized in opposite
10 polarity." Genomics 16:503-511 (1993).

Yamada, M. et al., "Preferential utilization of
specific immunoglobulin heavy chain diversity and
joining segments in adult human peripheral blood B
lymphocytes." J. Exp. Med. 173:395-407 (1991).

15

Equivalents

The foregoing description and Examples detail
certain preferred embodiments of the invention and
describes the best mode contemplated by the
inventors. It will be appreciated, however, that no
20 matter how detailed the foregoing may appear in text,
the invention may be practiced in many ways and the
invention should be construed in accordance with the
appended claims and any equivalents thereof.

- 95 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Abgenix, Inc.
- (ii) TITLE OF THE INVENTION: HUMAN MONOCLONAL ANTIBODIES TO EPIDERMAL GROWTH FACTOR RECEPTOR
- (iii) NUMBER OF SEQUENCES: 38
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: James F. Haley, FISH & NEAVE
 - (B) STREET: 1251 Ave. of the Americas
 - (C) CITY: New York
 - (D) STATE: New York
 - (E) COUNTRY: USA
 - (F) ZIP: 10020-1104
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: DOS
 - (D) SOFTWARE: FastSEQ Version 1.5
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 08/851,362
 - (B) FILING DATE: 05-MAY-1997
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Haley Jr., James F.
 - (B) REGISTRATION NUMBER: 27,794
 - (C) REFERENCE/DOCKET NUMBER: Cell 4.20 PCT
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (212) 596-9000
 - (B) TELEFAX: (212) 596-9090
 - (C) TELEX:

- 96 -

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CAGGTGCAGC TGGAGCAGTC GG

22

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GCTGAGGGAG TAGATCCTG AGGA

24

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 294 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

- 97 -

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GTCTCTGGTG	GCTCCATCAA	CAGTGGTGAT	TACTACTGGA	GCTGGATCCG	CCAGCACCCA	60
GGGAAGGGCC	TGGACTGCAT	TGGGTACATC	TATTACAGTG	GGAGCACCTA	CTACAACCCG	120
TCCCTCAAGA	GTCGAGTTAC	CATATCAGTA	GACACGTCTA	AGAATCAGTT	CTTCCTGAAG	180
CTGACCTCTG	TGACTGCCGC	GGACACGGCC	GTGTATTACT	GTGCGAGATC	TACGGTGGTA	240
AATCCGGGGT	GGTTCGACCC	CTGGGGCCAR	GGAACCCTGG	TCACCGTCTC	CTCA	294

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 264 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

ACCATCACTT	GCCAGGCGAG	TCAGGACATT	AACAACCTATT	TAAATTGGTT	TCAGCAGAAA	60
CCAGGGAAAG	CCCCTAAGGT	CCTGATCCAC	GATGCATCCA	ATTTGGAAAC	AGGGGGCCCA	120
TCAAGGTTCA	GTGGAAGTGG	ATCTGGGACA	GATTTTACTT	TCACCATCAG	CGGCCTGCAG	180
CCTGAAGACA	TTGCAACATA	TTATTGTCAA	CAGTATGAAA	GTCTCCCACT	CACTTTCGGG	240
GGAGGGACCA	AGGTGGGAT	CAAA				264

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 291 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GTCTCTGGTG	GCTCCATCAA	CAGTGGTGAT	TACTACTGGA	GCTGGATCCG	CCAGCACCCA	60
GGGAAGGGCC	TGGAGTGGAT	TGGGTCCATC	TATTACAGTG	GGAACACCTT	CTACAACCCG	120
TCCCTCAAGA	GTCGAGTTAC	CATATCACTA	GACACGTCTA	AGAACCAGTT	CTCCCTGAAG	180

- 98 -

CTGAGTTCTG TGACTGCCGC GGACACGGCC GTGTGTTACT GTGCGAGAAA TATAGTGACT 240
ACGGGTGCTT TTGATATCTG GGGCCAAGGG ACAATGGTCA CCGTCTCTTC A 291

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 264 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

ACCATCACTT GTCAGGCGAG TCAGGACATT ACCATTTATT TAAATTGGTA TCAACAGAAA 60
CCAGGGAAAG CCCCTAAGCT CCTGATCAAC GACGCATCCA GTTTGGAAAC AGGGGTCCCA 120
TTAAGGTTC GTGGAAGTGG ATCTGGGACA GATTTTACTT TCACCATCAG CAGCCTGCAG 180
CCTGAAGATA TTGCAACATA TTA CTGTGTCAG CAGTATGATC ATCTCCCGCT CACTTTCGGC 240
GCGGGGACCA AGGTGGCGAT CAAA 264

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 288 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GTCTCTGGTG GCTCCATCAG CAGTGGTGAT TACTACTGGA CCTGGATCCG CCAGCACCCA 60
GGGAAGGGCC TGGAGTGGAT TGGGTACATC TATTACAGTG GGAACACCTA CTACAACCCG 120
TCCCTCAAGA GTCGAGTTTC CATGTCAATA GACACGTCTG AGAACCAGTT CTCCTGAAG 180
CTGAGCTCTG TGACTGCCGC GGACACGGCC GTGTATTACT GTGCGAGAAA ACCAGTGACT 240
GGGGGGGAGG ACTACTGGGG CCAGGGAACC CTGGTCACCG TCTCCTCA 288

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- 99 -

- (A) LENGTH: 262 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

ACCATCACTT GCCAGGCGAG TCAGGACATT AGTAACTATT TAAATTGGTA TCAGCAGAAA	60
CCAGGGAAAG CCCTAAGCTC CTGATCTACG ATGCTTCCAA TTTGGAAACA GGGGTCCCAT	120
CAAGGTTCAG TGGAGTGGAT CTGGGACAGA TTTACTTTC ACCATCAGCA GCCTGCAGCC	180
TGAAGATGTT GGAACATATG TCTGTCAACA GTATGAGAGT CTCCCGTGGC GTTTTGGCCA	240
GGGGACCAAA CTGGAGATCA AA	262

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 291 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GTCTCTGGTG GCTCCATCAA CAGTGGTGAT TTCTACTGGA GCTGGATCCG CCAACACCCA	60
GGGAAGGGCC TGGAGTGGAT TGGGTACATC TATTACAGTG GGAGCACCTA CTACAACCCG	120
TCCCTCAAGA GTCGAGTTAC CATGTCAATA GACCCGTCTA AGAACCAGTT CTCCCTGAAA	180
CTGATCTCTG TGA CTGCCGC GGACACGGCC GTTTATTACT GTGCGACNTC CCTTTACTAT	240
GGCGGGGGTA TGGACGTCTG GGGCCAAGGG ACCACGGTCA CCGTCTCCTC A	291

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 264 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: cDNA

- 100 -

- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ACCATCACTT	GCCAGGCGAG	TCAGGACATT	AACAACTATT	TGAATTGGTA	TCAGCAGAGG	60
CCNGGGAACG	CCCCTAACT	CCTGATCTAC	GATGCATCCA	ATTTGGAAAC	AGGGGTCCCA	120
TCAAGGTTC	GTGGAAGTGG	ATCTGGGACA	GATTTTACTT	TCACCATCAA	CAGCCTGCAG	180
CCTGAAGATA	TTGCGACATA	TTATTGTCAA	CACTATGATC	ATCTCCCGTG	GACGTTCCGG	240
CAAGGGACCA	AGGTGGAANT	CAAA				264

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 291 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GTCTCTGGTG	GCTCCATCAA	CAATGGTGAT	TACTACTGGA	GCTGGATCCG	CCAGCACCCA	60
GGGAAGGGCC	TGGAGTGGAT	TGGGCACATC	TATTACAGTG	GGAGCACCTA	CTACATCCCG	120
TCCCTCAAGA	GTGGAAGTAC	CATATCAGTA	GACACGTCTA	AGAACCAGTT	CTCCCTGAAG	180
CTGAACTCTG	TGACTGCCGC	GGACACGGCC	GTGTATTACT	GTGCGAGAGG	GACAGTAACT	240
ACGTACTACT	TTGACTACTG	GGGCCAGGGA	ACCCTGGTCA	CCGTCTCCTC	A	291

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 270 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

- 101 -

ACCATCACTT	GCCGGGCAAG	TCAGAGCATT	AGCAGCTATT	TAAATTGGTA	TCAGCAGAAA	60
CCAGGGGAAAG	CCCCTAAGCT	CCTGATCTAT	GCTGCATCCA	GTTTGCAAAG	TGGGGTCCCA	120
TCAAGGTTCA	GTGGCAGTGG	ATCTGGGACA	GATTTCATCT	TCACCATCAG	CAGTCTGCAA	180
CCTGAAGATT	TTGCAACTTA	CTACTGTCAA	CAGGGTTACA	GAACCCCTCC	GGAGTGCACT	240
TTTGGCCAGG	GGACCAAGCT	GGAGATCAAA				270

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 291 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GTCTCTGGTG	GCTCCGTCAG	CAGTGGTGAT	TACTACTGGA	GCTGGATCCG	GCAGCCCCCA	60
GGGAAGGGAC	TGGAGTGGAT	TGGACATCTC	TATTACAGTG	GGAACACCAA	CTACAACCCC	120
TCCCTCAAGA	GTGAGTCAC	CATATCATT	GACACGTCCA	AGAACCAGTT	CTCCCTGAAG	180
CTGAGCTCTG	TGACCGCTGC	GGACACGGCC	GTGTATTACT	GTGCGAGAGA	TTTTTTGACT	240
GGTTCCTTCT	TTGACTACTG	GGGCCAGGGA	ACCCTGGTCA	CCGTCTCCTC	A	291

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 264 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

ACCATCACTT	GCCAGGCGAG	TCAGGACATA	AGCAACTATT	TAAATTGGTA	TCAGCAGAAA	60
CCAGGGGAAAG	CCCCTAAGCT	CCTGATCAAC	GATGCATCCG	ATTGGGAAAC	AGGGGTCCCA	120
TCAAGGATCA	GTGGAAGTGG	ATCTGGGACA	GATTTTACTT	TCACCATCAG	CAACCTGCAG	180
CCTGAAGATA	TTGCAACATA	TTACTGTCAA	CAATATGATA	GTCTCCCGCT	CACTTTCGGC	240
GGAGGGACCA	AGGTGGAGAT	CAGA				264

(2) INFORMATION FOR SEQ ID NO:15:

- 102 -

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 288 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GTCTCTGGTG	GCTCCGCTCTA	CAGTGGTGAT	TACTACTGGA	GCTGGATCCG	GCAGCCCCCC	60
GGGAAGGGAC	TGGAGTGGAT	TGGGTATATC	TATTACAGTG	GGAGCACCAA	TTACAATCCC	120
TCCCTCAAGA	GTCGAGTCAC	CATATCAGTA	GACAGGTCCA	AGAACCAGTT	CTCCCTGAAG	180
CTGAGCTCTG	TGACCGCTGC	GGACACGGCC	GTGTATTACT	GTGCGAGAGA	CTCCACTACTG	240
GGAGCTACCA	ACTACTGGGG	CCAGGGAACC	CTGGTCACCG	TCTCCTCA		288

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 264 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

ACCATCACTT	GCCAGGCGAG	TCNGGACATT	AATAACTATT	TANATTGGTN	TCAGCAGAAA	60
CCAGGGAAG	CCCCTAAAT	CCTGATCTCC	GATGCATCCA	ATTTAGAAAC	AGGGGTCCCA	120
TCGAGGTTCA	GTGGAAGTGG	ATCTGGGACA	GANTNTACTT	TCACCATCAG	CAGCCTGCAG	180
CCTGAAGATA	TTGCNACATA	TCACTGTCNA	CAGTATNATA	GTCTCCCGCT	CACTTTCGGC	240
GGAGGGACCA	AGGTAGAGAT	CAAA				264

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 288 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- 103 -

- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GTCTCTGGTG	GCTCCGTCAG	CAGTGGTGAT	TACTACTGGA	CCTGGATCCG	GCAGTCCCCA	60
GGGAAGGGAC	TGGAGTGGAT	TGGACACATC	TATTACAGTG	GGAACACCAA	TTATAACCCC	120
TCCCTCAAGA	GTGACTCAC	CATATCAATT	GACACGTCCA	AGACTCAGTT	CTCCCTGAAG	180
CTGAGTTCTG	TGACCGCTGC	GGACACGGCC	ATTTATTACT	GTGTGCGAGA	TCGAGTGACT	240
GGTGCTTTTG	ATATCTGGGG	CCAAGGGACA	ATGGTCACCG	TCTCTTCA		288

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 264 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

ACCATCACTT	GCCAGGCGAG	TCAGGACATC	AGCAACTATT	TAAATGGTA	TCAGCAGAAA	60
CCAGGGAAAG	CCCCTAAACT	CCTGATCTAC	GATGCATCCA	ATTTGGAAC	AGGGGTCCCA	120
TCAAGGTTCA	GTGGAAGTGG	ATCTGGGACA	GATTTTACTT	TCACCATCAG	CAGCCTGCAG	180
CCTGAAGATA	TTGCAACATA	TTTCTGTCAA	CACTTTGATC	ATCTCCCGCT	CGCTTTCGGC	240
GGAGGGACCA	AGGTGGAGAT	CAAA				264

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 76 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE: internal
- (vi) ORIGINAL SOURCE:

- 104 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

```

Val Ser Gly Gly Ser Ile Ser Ser Gly Gly Tyr Tyr Trp Ser Trp Ile
 1           5           10           15
Arg Gln His Pro Gly Lys Gly Leu Glu Trp Ile Gly Tyr Ile Tyr Tyr
 20           25           30
Ser Gly Ser Thr Tyr Tyr Asn Pro Ser Leu Lys Ser Arg Val Thr Ile
 35           40           45
Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu Lys Leu Ser Ser Val
 50           55           60
Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg
 65           70           75

```

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 76 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

```

Thr Ile Thr Cys Gln Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn Trp
 1           5           10           15
Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Asp Ala
 20           25           30
Ser Asn Leu Glu Thr Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser
 35           40           45
Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Leu Gln Pro Glu Asp Ile
 50           55           60
Ala Thr Tyr Tyr Cys Gln Gln Tyr Asp Asn Leu Pro
 65           70           75

```

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 76 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

- 105 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

```

Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser Ser Tyr Leu Asn Trp
 1           5           10           15
Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Ala Ala
      20           25           30
Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser
 35           40           45
Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe
 50           55           60
Ala Thr Tyr Tyr Cys Gln Ser Tyr Ser Thr Pro
65           70           75

```

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 76 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

```

Val Ser Gly Gly Ser Val Ser Ser Gly Ser Tyr Tyr Trp Ser Trp Ile
 1           5           10           15
Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile Gly Tyr Ile Tyr Tyr
      20           25           30
Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys Ser Arg Val Thr Ile
 35           40           45
Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu Lys Leu Ser Ser Val
 50           55           60
Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg
65           70           75

```

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 98 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

- 106 -

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

```

Val Ser Gly Gly Ser Ile Asn Ser Gly Asp Tyr Tyr Trp Ser Trp Ile
 1           5           10           15
Arg Gln His Pro Gly Lys Gly Leu Asp Cys Ile Gly Tyr Ile Tyr Tyr
      20           25           30
Ser Gly Ser Thr Tyr Tyr Asn Pro Ser Leu Lys Ser Arg Val Thr Ile
 35           40           45
Ser Val Asp Thr Ser Lys Asn Gln Phe Phe Leu Lys Leu Thr Ser Val
 50           55           60
Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg Ser Thr Val Val
 65           70           75           80
Asn Pro Gly Trp Phe Asp Pro Trp Gly Gln Gly Thr Leu Val Thr Val
      85           90           95
Ser Ser

```

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 105 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

```

Thr Ile Thr Cys Gln Ala Ser Gln Asp Ile Asn Asn Tyr Leu Asn Trp
 1           5           10           15
Phe Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile His Asp Ala
      20           25           30
Ser Asn Leu Glu Thr Gly Gly Pro Ser Arg Phe Ser Gly Ser Gly Ser
 35           40           45
Gly Thr Asp Phe Thr Phe Thr Ile Ser Gly Leu Gln Pro Glu Asp Ile
 50           55           60
Ala Thr Tyr Tyr Cys Gln Gln Tyr Glu Ser Leu Pro Leu Thr Phe Gly
 65           70           75           80
Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val
      85           90           95
Phe Ile Phe Pro Pro Ser Asp Glu Gln
      100           105

```

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 97 amino acids

- 107 -

(B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
 (iii) HYPOTHETICAL: NO
 (iv) ANTISENSE: NO
 (v) FRAGMENT TYPE: internal
 (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

```

Val Ser Gly Gly Ser Ile Asn Ser Gly Asp Tyr Tyr Trp Ser Trp Ile
 1           5           10           15
Arg Gln His Pro Gly Lys Gly Leu Glu Trp Ile Gly Ser Ile Tyr Tyr
          20           25           30
Ser Gly Asn Thr Phe Tyr Asn Pro Ser Leu Lys Ser Arg Val Thr Ile
          35           40           45
Ser Leu Asp Thr Ser Lys Asn Gln Phe Ser Leu Lys Leu Ser Ser Val
          50           55           60
Thr Ala Ala Asp Thr Ala Val Cys Tyr Cys Ala Arg Asn Ile Val Thr
          65           70           75           80
Thr Gly Ala Phe Asp Ile Trp Gly Gln Gly Thr Met Val Thr Val Ser
          85           90           95
Ser
  
```

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 105 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
 (iii) HYPOTHETICAL: NO
 (iv) ANTISENSE: NO
 (v) FRAGMENT TYPE: internal
 (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

```

Thr Ile Thr Cys Gln Ala Ser Gln Asp Ile Thr Ile Tyr Leu Asn Trp
 1           5           10           15
Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Asn Asp Ala
          20           25           30
Ser Ser Leu Glu Thr Gly Val Pro Leu Arg Phe Ser Gly Ser Gly Ser
          35           40           45
Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Leu Gln Pro Glu Asp Ile
          50           55           60
Ala Thr Tyr Tyr Cys Gln Gln Tyr Asp His Leu Pro Leu Thr Phe Gly
          65           70           75           80
Gly Gly Thr Lys Val Ala Ile Lys Arg Thr Val Ala Ala Pro Ser Val
  
```

- 108 -

Phe Ile Phe Pro Pro Ser Asp Glu Gln 85 90 95
100 105

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 96 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Val Ser Gly Gly Ser Ile Ser Ser Gly Asp Tyr Tyr Trp Thr Trp Ile
1 5 10 15
Arg Gln His Pro Gly Lys Gly Leu Glu Trp Ile Gly Tyr Ile Tyr Tyr
20 25 30
Ser Gly Asn Thr Tyr Tyr Asn Pro Ser Leu Lys Ser Arg Val Ser Met
35 40 45
Ser Ile Asp Thr Ser Glu Asn Gln Phe Ser Leu Lys Leu Ser Ser Val
50 55 60
Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg Lys Pro Val Thr
65 70 75 80
Gly Gly Glu Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
85 90 95

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 105 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Thr Ile Thr Cys Gln Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn Trp
1 5 10 15
Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Asp Ala

- 109 -

20							25				30				
Ser	Asn	Leu	Glu	Thr	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser
35							40				45				
Gly	Thr	Asp	Phe	Thr	Phe	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Glu	Asp	Ile
50							55				60				
Val	Gly	Tyr	Tyr	Val	Gln	Gln	Tyr	Glu	Ser	Leu	Pro	Cys	Gly	Phe	Gly
65							70				75				
Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val
85							90				95				
Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln							
100							105								

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 97 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEO ID NO:29:

[illegible]

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 105 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO

- 110 -

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

```

Thr Ile Thr Cys Gln Ala Ser Gln Asp Ile Ser Asn Asn Leu Asn Trp
 1           5           10           15
Tyr Gln Gln Lys Arg Gly Asn Ala Pro Lys Leu Leu Ile Tyr Asp Ala
      20           25           30
Ser Asn Leu Glu Thr Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser
      35           40           45
Gly Thr Asp Phe Thr Phe Thr Ile Ser Asn Leu Gln Pro Glu Asp Ile
      50           55           60
Ala Thr Tyr Tyr Cys Gln His Tyr Asp His Leu Pro Trp Thr Phe Gly
      65           70           75           80
Gln Gly Thr Lys Val Glu Xaa Lys Arg Thr Val Ala Ala Pro Ser Val
      85           90           95
Phe Ile Phe Pro Pro Ser Asp Glu Gln
      100           105

```

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 97 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

```

Val Ser Gly Gly Ser Ile Asn Asn Gly Asp Tyr Tyr Trp Ser Trp Ile
 1           5           10           15
Arg Gln His Pro Gly Lys Gly Leu Glu Trp Ile Gly His Ile Tyr Tyr
      20           25           30
Ser Gly Ser Thr Tyr Tyr Ile Pro Ser Leu Lys Ser Arg Thr Thr Ile
      35           40           45
Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu Lys Leu Asn Ser Val
      50           55           60
Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg Gly Thr Val Thr
      65           70           75           80
Thr Tyr Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser
      85           90           95
Ser

```

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- 111 -

- (A) LENGTH: 107 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE: internal
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

```

Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser Ser Tyr Leu Asn Trp
 1           5           10           15
Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Ala Ala
 20           25           30
Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser
 35           40           45
Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe
 50           55           60
Ala Thr Tyr Tyr Cys Gln Gln Gly Tyr Arg Thr Pro Pro Glu Cys Ser
 65           70           75           80
Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala Pro
 85           90           95
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
 100           105

```

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 97 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

```

Val Ser Gly Gly Ser Val Ser Ser Gly Asp Tyr Tyr Trp Ser Trp Ile
 1           5           10           15
Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile Gly His Leu Tyr Tyr
 20           25           30
Ser Gly Asn Thr Asn Tyr Asn Pro Ser Leu Lys Ser Arg Val Thr Ile
 35           40           45
Ser Leu Asp Thr Ser Lys Asn Gln Phe Ser Leu Lys Leu Ser Ser Val
 50           55           60
Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg Asp Phe Leu Thr

```

65 70 75 80
Gly Ser Phe Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser .
 85 90 95
Ser

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

- 113 -

```

Val Ser Gly Gly Ser Val Tyr Ser Gly Asp Tyr Tyr Trp Ser Trp Ile
 1           5           10           15
Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile Gly Tyr Ile Tyr Tyr
 20           25           30
Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys Ser Arg Val Thr Ile
 35           40           45
Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu Lys Leu Ser Ser Val
 50           55           60
Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg Asp Ser Ile Leu
 65           70           75           80
Gly Ala Thr Asn Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 85           90           95

```

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 105 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

```

Thr Ile Thr Cys Gln Ala Ser Gln Xaa Ile Ser Asn Tyr Leu Xaa Trp
 1           5           10           15
Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Xaa Leu Ile Ser Asp Ala
 20           25           30
Ser Asn Leu Glu Thr Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser
 35           40           45
Gly Thr Xaa Xaa Thr Phe Thr Ile Ser Ser Leu Gln Pro Glu Asp Ile
 50           55           60
Ala Thr Tyr His Cys Xaa Gln Tyr Xaa Ser Leu Pro Leu Thr Phe Gly
 65           70           75           80
Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val
 85           90           95
Phe Ile Phe Pro Pro Ser Asp Glu Gln
 100           105

```

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 95 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

-114-

- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE: internal
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

```

Val Ser Gly Gly Ser Val Ser Ser Gly Asp Tyr Tyr Trp Thr Trp Ile
 1           5           10           15
Arg Gln Ser Pro Gly Lys Gly Leu Glu Trp Ile Gly His Ile Tyr Tyr
 20           25           30
Ser Gly Asn Thr Asn Tyr Asn Pro Ser Leu Lys Ser Arg Leu Thr Ile
 35           40           45
Ser Ile Asp Thr Ser Lys Thr Gln Phe Ser Leu Lys Leu Ser Ser Val
 50           55           60
Thr Ala Ala Asp Thr Ala Ile Tyr Tyr Cys Val Arg Asp Arg Val Thr
 65           70           75           80
Gly Ala Phe Asp Ile Trp Gly Gln Gly Thr Met Val Thr Ser Ser
 85           90           95

```

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 105 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE: internal
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

```

Thr Ile Thr Cys Gln Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn Trp
 1           5           10           15
Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Asp Ala
 20           25           30
Ser Asn Leu Glu Thr Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser
 35           40           45
Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Leu Gln Pro Glu Asp Ile
 50           55           60
Ala Thr Tyr Phe Cys Gln His Phe Asp His Leu Pro Leu Ala Phe Gly
 65           70           75           80
Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val
 85           90           95
Phe Ile Phe Pro Pro Ser Asp Glu Gln
100           105

```


- 115 -

ClaimsWhat We Claim Is:

1. An antibody against epidermal growth factor receptor comprising a heavy chain variable region amino acid sequence wherein a portion of the sequence is encoded by a human V_H 4 family gene and any of the mutations thereto represented by the nucleotide sequences shown in Figures 2, 6, 10, 14, 18, 22, 26, and 30.

2. The antibody of Claim 1, wherein the heavy chain variable region amino acid sequence comprises an Aspartic Acid amino acid substitution at residue 10.

3. An antibody against epidermal growth factor receptor comprising a heavy chain variable region amino acid sequence wherein a portion of the sequence is encoded by a human V_H 4-31 gene and any of the mutations thereto represented by the nucleotide sequences shown in Figures 2, 6, 10, 14, and 18.

4. The antibody of Claim 3, wherein the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:23.

5. The antibody of Claim 4, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:24.

- 116 -

6. The antibody of Claim 3, wherein the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:25.

7. The antibody of Claim 6, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:26.

8. The antibody of Claim 3, wherein the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:27.

9. The antibody of Claim 8, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:28.

10. The antibody of Claim 3, wherein the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:29.

11. The antibody of Claim 10, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:30.

12. The antibody of Claim 3, wherein the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:31.

-117-

13. The antibody of Claim 12, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:32.

14. An antibody against epidermal growth factor receptor comprising a heavy chain variable region amino acid sequence wherein a portion of the sequence is encoded by a human V_H 4-61 gene and any of the mutations thereto represented by the nucleotide sequences shown in Figures 22, 26, and 30.

15. The antibody of Claim 14, wherein the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:33.

16. The antibody of Claim 15, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:34.

17. The antibody of Claim 14, wherein the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:35.

18. The antibody of Claim 17, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:36.

19. The antibody of Claim 14, wherein the heavy chain variable region comprises the contiguous sequence from CDR1 through CDR3 as represented in SEQ

- 118 -

ID NO:37.

20. The antibody of Claim 19, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:38.

21. An antibody against epidermal growth factor receptor comprising a light chain variable region amino acid sequence wherein a portion of the sequence is encoded by a human Vk I family gene and any of the mutations thereto represented by the nucleotide sequences shown in Figures 4, 8, 12, 16, 20, 24, 28, and 32.

22. The antibody of Claim 21, wherein the light chain variable region comprises the sequence represented by SEQ ID NO:24.

23. The antibody of Claim 21, wherein the light chain variable region comprises the sequence represented by SEQ ID NO:26.

24. The antibody of Claim 21, wherein the light chain variable region comprises the sequence represented by SEQ ID NO:28.

25. The antibody of Claim 21, wherein the light chain variable region comprises the sequence represented by SEQ ID NO:30.

26. The antibody of Claim 21, wherein the light chain variable region comprises the sequence

- 119 -

represented by SEQ ID NO:32.

27. The antibody of Claim 21, wherein the light chain variable region comprises the sequence represented by SEQ ID NO:34.

28. The antibody of Claim 21, wherein the light chain variable region comprises the sequence represented by SEQ ID NO:36.

29. The antibody of Claim 21, wherein the light chain variable region comprises the sequence represented by SEQ ID NO:38.

30. An antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:23.

31. The antibody of Claim 30, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:24.

32. An antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:25.

33. The antibody of Claim 32, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:26.

- 120 -

34. An antibody against epidermal growth factor receptor comprising a heavy chain variable region comprises a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:27.

35. The antibody of Claim 34, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:28.

36. An antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:29.

37. The antibody of Claim 36, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:30.

38. An antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:31.

39. The antibody of Claim 38, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:32.

40. An antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:33.

- 121 -

41. The antibody of Claim 40, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:34.

42. An antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:35.

43. The antibody of Claim 42, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:36.

44. An antibody against epidermal growth factor receptor comprising a heavy chain variable region comprising a contiguous sequence from CDR1 through CDR3 as represented in SEQ ID NO:37.

45. The antibody of Claim 44, further comprising a light chain variable region comprising the sequence represented by SEQ ID NO:38.

46. In a method for treating a solid tumor with an antibody against epidermal growth factor receptor, the improvement comprising administering to a patient having a solid tumor an antibody of any one of claims 1-45.

VS⁶⁶SI¹N⁵⁶SG¹DYYWS¹IRQHPKGL¹DCIG¹YIYSGSTY¹NP¹SLKSRVTI¹SVDT¹SKNQF¹FLKL¹TSVTAADIAVYCARSTV¹NP¹GVFDPVGGGLVT¹VSS (SEQ ID NO: 23)

CDR1 CDR2 CDR3

FIG. 1

GTCTCTGGTG GCTCCATCAA CAGTGGTGAT TACTACTGGA GCTGGATCCG CCAGCACCCA GGGAGGGCC TGGACTGCAT TGGGTACATC TATTACAGTG GGAGCACCTA CTACAACCCG
TCCCTCAAG GTGAGTTAC CATACAGTA GACAGTCTA AGATCAGTT CTCTGAAG CTGACTCTCG TGACTGGCCG GGACAGGCC GTGTATTACT GTGCGAGATC TACGGTGGTA
AATCCGGGGT GGTTCGACCC CTGGGGCCAR GGAACCCCTG TCACCGTCTC CTCA (SEQ ID NO: 3)

FIG. 2

TI¹TQASQDI¹N¹N¹YL¹N¹WF¹QKPKGAPK¹VLI¹HDASNLETG¹GPSRFSGSGSDFTFTIS¹GLQPEDIA¹TYCQRY¹ESLPLIFGGGK¹VEIKRTVAAPSVFI¹PPSDEQ (SEQ ID NO: 24)

CDR1 CDR2 CDR3

FIG. 3

ACCATCACTT GCCAGGGGAG TCAGGACATT AACAACTATT TAAATTGGTT TCAGCAGAAA CCAGGGAAG CCCCT AAGGTCCTGA TCCACGATGC ATCCAATTG GAAACAGGGG
GCCATCAAG GTTCAGTGA AGTGGATCTG GGACA-GATTTTACTT TCACCATCAG CGGCTGCAG CCTGAAGACA TTGCACATA TTATTGCAA CAGTATGAA GTCTC CCACTCACTT
TCGGCGGAGG GACCAAGGTG GAGATCAAA (SEQ ID NO: 4)

FIG. 4

1/20

2/20

VSGGSI NsgDYYWSVIRHPGKLENI GSIYYSG NtFYNPSLKSRTVITSLDTSKNQFSLKLSSTVADTAVCYCARNIVTTGAFDI VGGGTMTVSS (SEQ ID NO: 25)
 CDR1 CDR2 CDR3

FIG. 5

GTCTCTGGTG GCTCCATCAA CAGTGGTGAT TACTACTGGA GCTGGATCCG CCAGCACCCA GGAAGGGCC TGGAGTGGAT TGGGTCCATC TATTACAGTG GGAACACCTT CTACACCCG
 TCCCTCAAGA GTCGAGTTAC CATATCACTA GACAGCTCTA AGAACCACTT CTCCTGAG CTGAGTTCTG TGACTGCCG GACACGGCC GTGTGTACT GTGGAGAAA TATAGTACT
 ACGGGTGCTT TTGATATCTG GGGCCAAGG ACAATGGTCA CCGTCTCTC A (SEQ ID NO: 5)

FIG. 6

TITCDASQDIT IYLVYQKPKAPKLLI NdasSLETGVPLRFSGSGSDTFTITSSLPEDIATYYCQDYHLPLTFGGGTVAKRTVAAPSVFIFPPSDEQ (SEQ ID NO: 26)
 CDR1 CDR2 CDR3

FIG. 7

ACCATCACTT GTCAGGCGAG TCAGGACATT ACCATTATT TAAATTGGTA TCACAGAAA CCAGGGAAG CCCT AGCTCCTGA TCAACACGC ATCCAGTTG GAAACAGGGG
 TCCATTAAAG GTTCAGTGA AGTGGATCTG GGACA GATTITACTT TCACCATCAG CAGCCTGCAG CTTGAAGATA TTGAACATA TTAGTGCAA CAGTATGATC ATCTC CCGCTCACTT
 TCGGCGGGCG GACCAAGGTG GCGATCAAA (SEQ ID NO: 6)

FIG. 8

3/20

VSGGSISSGDYYVTWIRQHPKGLWIGYIYSGNTYINPSLKSRVSMsIDtsENDFSLKSSVTAADTAVYCARPKVITGGEDYGGGTLVTYSS (SEQ ID NO: 27)

CDR1

CDR2

CDR3

FIG. 9

GTCTCTGGTG GCTCCATCAG CAGTGGTGAT TACTACTGGA CTGGATCCG CCAGCACCCA GGGAGGGCC TGGGTACATC TATTACAGTG GGAACACCTA CTACAACCCG
TCCCTCAAGA GTGAGTTTC CATGTCAATA GACAGTCTG AGAACCAAGT CTCCCTGAAG CTGAGCTCTG TGACTGCCGC GGACACGGCC GTGTATTACT GTGCAGAGAA ACCAGTGACT
GGGGGGGAGG ACTACTGGGG CCAGGGAACC CTGGTCACCG TCTCCTCA (SEQ ID NO: 7)

FIG. 10

TTTCAASQDI SNYLNWYQKPKAPKLLI YDASNLETGVPSPRSGSGSDTFITISSLPEDI VGYVQDY^{ES}LP^{CG}FGG^{GT}KL^{ET}KRTVAAPSVFI^{PP}SD^{EQ} (SEQ ID NO: 28)

CDR1

CDR2

CDR3

FIG. 11

ACCATCATT GCCAGGGGAG TCAGGACATT AGTAACATT TAAATTGGTA TCAGCAGAAA CCAGGGAAG CCT AAGCTCCTGA TCTAGGATGC TTCCAATTG GAAACAGGGG TCCCATCAAG
GTTCAAGTGA GTGATCTG GGACA GATTTACTT TCACATCAG CAGCCTGCAG CCTGAAGATG TTGGAACATA TGTCTGTCAA CAGTATGAGA GTCTC CCGTGGCGTT TTGGCCAGGG
GACCAAACTG GAGATCAAA (SEQ ID NO: 8)

FIG. 12

4/20

VSGGSI Nsg DF YNSVIRDPKGLKLEVI GYIYYSGSTYNNPSLKSRVT Ms In P SKNQFSKL ISVTAADIAYYCA T SLYYGGMDYVGGGTTVTVSS (SEQ ID NO: 29)
 CDR1 CDR2 CDR3

FIG. 13

GTCTCTGGTG GCTCCATCAA CAGTGGTGAT TTCTACTGGA GCTGGATCCG CCAACACCCA GGAAGGGCC TGGAGTGGAT TGGGTACATC TATTACAGTG GGAGCACCTA CTACACCCG
 TCCCTCAAGA GTGAGATTAC CATGTCAATA GACCGTCTA AGAACCAATT CTCCCTGAAA CTGATCTCTG TGACTGCCGC GGACACGGCC GTTATTACT GTGCGACNTC CCTTACTAT
 GGGGGGGGTA TGGACGTCTG GGGCCCAAGG ACCACGGTCA CCGTCTCCTC A (SEQ ID NO: 9)

FIG. 14

TLTCCASDJSN LNwYQOK Rg NApKLLIYDASNLETGVP SRFSGSGSDTFTTISNLPEDIATYYCQ H YD LHPVTEGGGTVKXRTVAAPSVFIPPSDEQ (SEQ ID NO: 30)
 CDR1 CDR2 CDR3

FIG. 15

ACCATCATT GCCAGGCGAG TCAGGACATT AACAACTATT TGAATTGGTA TCAGCAGAGG CCGGGGAACG CCCCT AACCTCCTGA TCTAGGATGC ATCCAATTG GAACACGGGG
 TCCCATCAAG GTTCAGTGA AGTGGATCTG GGACA GATTITACTT TCACCATCAA CAGCCTGCAG CCTGAAGATA TTGGACATA TTATTGTCAA CACTATGATC ATCTC CCGTGGACGT
 TCGGCCAAGG GACCAAGGTG GAANTCAAA (SEQ ID NO: 10)

FIG. 16

5/20

VSGGSINNGDYYSWIRQHPKGLWIGHIYYSGSTYYIPSLKSR
 T T I S V D T S K N Q F S L K L N S V T A A D I A V Y Y C A R G T V I T Y Y F D Y W G Q G T T V T V S S (SEQ ID NO: 31)
 CDR1 CDR2 CDR3

FIG. 17

GTCTCTGGTG GCTCCATCAA CAATGGTGAT TACTACTGGA GCTGGATCCG CCAGCACCCA GGGAGGGCC TGGAGTGGAT TGGGCACATC TATTACAGTG GGAGCACCTA CTACATCCCC
 TCCCTCAAGA GTCGAACTAC CATATCAGTA GACAGTCTA AGAACCAGTT CTCCTGAAG CTGAACCTG TGACTGCCG GGACACGCCC GTGTATTACT GTGCGAGAGG GACAGTAACT
 ACGTACTACT TTGACTACTG GGGCCAGGGA ACCCTGGTCA CCGTCTCCTC A (SEQ ID NO: 11)

FIG. 18

T I T C R A S Q S I S S Y L N W Y Q K P G K A P K L L I Y A A S S L O S G V P S R F S G S G S G T D F T L T I S S L O P E D F A T Y Y C Q Q G Y R T P P E C S F G G G K L E I K R T V A P S V F I F P P S D E Q (SEQ ID NO: 32)
 CDR1 CDR2 CDR3

FIG. 19

ACCATCACTT GCCGGGCAAG TCAGAGCATT AGCAGCTATT TAAATTGGTA TCAGCAGAAA CCAGGGAAG CCCCT AAGCTCTGA TCTATGCTGC ATCCAGTTTG CAAAGTGGG
 TCCCATCAAG GTTCAGTGGC AGTGGATCTG GGACA GATTTCATC TCACCATCAG CAGCTGCAA CCTGAAGATT TTGCAACTTA CTACGTGCAA CAGGGTTACA GAACC CCTCCGGAGT
 GCAGTTTTTG CCAGGGGACC AAGCTGGAGA TCAAA (SEQ ID NO: 12)

FIG. 20

6/20

VSGSVSSG DYYKSHIRQPPGKGLVIGHL YYSGNTNYPNPKSRVTISL DTSKNQFSLKSSVTAADTAVYYCARDL TGSFFDYVGGTILVTSS (SEQ ID NO: 33)

CDR1 CDR2 CDR3

FIG. 21

GTCTCTGGTG GCTCCGTCAG CAGTGGTGAT TACTACTGGA GCTGGATCCG GCAGCCCCCA GGAAGGGAC TGGAGTGGAT TGGACATCTC TATTACAGTG GGAACACCAA CTACAACCC

TCCCTCAAGA GTCCAGTCAC CATATGATTA GACAGTCCA AGAACAGATT CTCCCTGAAG CTGAGCTCTG TGACCGCTGC GCACACGGCC GTGTATTACT GTCCGAGAGA TTTTGTGACT

GGTTCTTCT TTGACTACTG GGGCCAGGGA ACCCTGGTCA CCGTCTCCTC A (SEQ ID NO: 13)

FIG. 22

TI TCQASQDI SNYLNWYQQKPKAPKLLI NDASD LETGVP SRISGSGGIDFIFTIS NLQPEDIAIYYCQQYD SLPLIFGGGKVEIRTVAPSVFI PPPSDEQ (SEQ ID NO: 34)

CDR1 CDR2 CDR3

FIG. 23

ACCATCACTT GCCAGCGGAG TCAGGACATA AGCAACTATT TAAATTGGTA TCAGCAGAAA CCAGGGAAG CCCCT AAGCTCCTGA TCACGATGC ATCCGATTG GAAACAGGGG

TCCCATCAAG GATCAGTGA AGTGGATCTG GGACA GATTITACT TCACCATCAG CAACCTGCAG CCTGAAGATA TTCAACATA TTACTGTCAA CAATATGATA GTCTC CCGCTCACTT

TCGGCGGAGG GACCAAGGTG GAGATCAGA (SEQ ID NO: 14)

FIG. 24

7/20

VSGGSV YsgDyywswrpppgkglewlgtyyysgsnynpslksrvttsvdtsknqfslklssvtaadtavyyccardsilgatnyhgggtlvtvss (SEQ ID NO: 35)

CDR1

CDR2

CDR3

FIG. 25

GTCTCTGGTG GCTCCGTCTA CAGTGGTGAT TACTACTGGA GCTGGATCCG GCAGCCCCC GGAAGGGAC TGGATTGGAT TGGGTATATC TATTACAGTG GGAGCACCAA TTACAATCCC
TCCCTCAAGA GTGAGTCAC CATATCAGTA GACAGTCCA AGAACCAAGT CTCCTGAAG CTGAGCTCTG TGACCGCTGC GGACACGGCC GTGTATTACT GTGGGAGAGA CTCCTACTG
GGAGCTACCA ACTACTGGGG CCAGGGAACC CTGGTCACCG TCTCCTCA (SEQ ID NO: 15)

FIG. 26

TI TCQASQX₁ISNYL XuyqkpgkapkX_{LI}S_DASNLEITGVPSRFSGSGSGTXX₁TFITSSLPEDIATYHcX_{QY}X_{SL}PLTFGGGKVEIKRTVAAPSVFIPPSDEQ (SEQ ID NO: 36)

CDR1

CDR2

CDR3

FIG. 27

ACCATCACTT GCCAGGCGAG TCNGGACATT AATACTATT TANATTGGTN TCACGAGAA CCAGGGAAAG CCCCT AAATCCTGA TCCCGATGC ATCCAATTTA GAAACAGGGG
TCCCATCGAG GTTCAGTGA AGTGGATCTG GGACA GANTNTACTT TCACCATCAG CAGCCTGCAG CCTGAAGATA TTGCNACATA TCACTGTGNA CAGTATNATA GTCTC CCGCTCACTT
TCGGCGGAGG GACCAAGGTA GAGATCAAA (SEQ ID NO: 16)

FIG. 28

VSGGSVSSGDYYVTWIRSPQKGLEWIGHIYYSGNINYNPSLKSRLTISIDTSKTQFSLKLSSTVAADTAIYYCVRDRVTGAFDINGGTHVTSS (SEQ ID NO: 37)

CDR1

CDR2

CDR3

FIG. 29

GTCTCTGGTG GCTCCGTGAG CAGTGGTGAT TACTACTGGA CCTGGATCCG GCAGTCCCA GGAAGGGAC TGGAGTGGAT TGGACACATC TATTACAGTG GGAACACCAA TTATAACCCC
TCCCTCAAGA GTGACTCAC CATATCAATT GACACGTCCA AGACTCAGTT CTCCTGAAG CTGAGTTCTG TGACCGCTGC GGACACGGCC ATTATTACT GTGTGCGAGA TCGAGTGACT
GGTGCTTTTG ATATCTGGGG CCAAGGGACAATGGTCACCG TCTCTTCA (SEQ ID NO: 17)

8/20

FIG. 30

TI TCQASQDI SNYLNWYQKPKGAPKLLI YDASNLETGVPSRFSGSGTDFTFTISSLPEDIATYFCQHF D H L P L A F G G G T K V E I K R T V A A P S V F I P P S D E Q (SEQ ID NO: 38)

CDR1

CDR2

CDR3

FIG. 31

ACCATCACTT GCCAGGGCAG TCAGGACATC AGCACTATT TAAATTGGTA TCAGCAGAAA CCAGGGAAG CCCCT AACTCCTGA TCTACGATGC ATCCAATTG GAAACAGGGG
TCCCATCAAG GTTCAGTGA AGTGGATCTG GGACA GATTTTACTT TCACCATCAG CAGCCTGCAG CCTGAAGATA TTTCGTGCAA CACTTTGATC AICTC CCGCTCGCTT
TCGGCGGAGG GACCAAGGTG GAGATCAAA (SEQ ID NO: 18)

FIG. 32

*Amino Acid Sequences and Structure of Human Heavy
Chain Derived from EGFR-Specific Hybridomas*

9/20

Human $\gamma 2$	CDR1	CDR2	CDR3	
4-31	VSGGSISSGGYYWSIRQHPGKGLEWIGYIYYSGSTYNPCLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCAR			ASTKGPSVFPPPLAPCSRSTST
E1.1	N-D	DC	STVWPG WFDP	WGQGLVTYSS
E2.4	N-D	S N-F	NIVTTG AFDI	WGQGTWTVSS
E2.5	D-T	N	KPVTGG EDY	WGQGLVTYSS
E6.2	N-DI	M-I-P	TSLYYGG GMDV	WGQGTWTVSS
E6.4	NN-D	I	GTVTYY YFDY	WGQGTWTVSS

	CDR1	CDR2	CDR3	
4-61	VSGGSVSSGGYYWSIRQPPGKGLEWIGYIYYSGSTYNPCLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCAR			
E2.11	D	HL N	DFLTGSF FDY	WGQGTWTVSS
E6.3	Y-D		DSILGA TNY	WGQGLVTYSS
E7.6.3	D-T	S L-I	V-DRVTGA FDI	WGQGTWTVSS

FIG. 33

*Amino Acid Sequence and Structure of Human Kappa
Chain Derived from EGFR-Specific Hybridomas*

	V _K SEGMENT			J _K SEGMENT		
	CDR1	CDR2	CDR3	CDR3	CDR3	CDR3
V _K (018)	TI TQASQDI SNV LNWYQQKPGKAPKLLIYDASNLETGVP	SRFSGSGGTDFTITLSSLPEDIAITYYQQQYDNL	P	RTVAAPSVFIFPPSDEQ		
Human Ck	---	---	---	---	---	---
EGF.1.1.K	N--F--N--	V--H--G--	ES--	LTGGGKVEIK	---	---
EGF.2.4.K	TI--	N--S--L--	H--	LTGGGKVAIK	---	---
EGF.2.5.K	---	---	VG--V--ES--	CGFGGKLEIK	---	---
EGF.2.11.K	---	N--D--I--	S--	LTGGGKVEIR	---	---
E6.2.K	N--R--N--	---	H--H--	WTFGGGKVEKK	---	---
E6.3.K	X--N--X--	X--S--XX--	H--X--XS--	LTGGGKVEIK	---	---
E6.4.K*	R--S--S--	A--S--QS--	F--	CSFGGKLEIK	---	---
*012 Gene	---	---	---	---	---	---
E7.6.3.K	---	---	F--HF--H--	LAFGGGKVEIK	---	---

10/20

FIG. 34

11/20




FIG. 35

12/20





FIG. 36

13/20

*Inhibition of TGF- α binding to A431 cells
by anti-EGF receptor antibodies*

**FIG. 37**

14/20

ABX-EGF: Blockage of EGF Binding to Human
Colon Adenocarcinoma SW948 Cells




FIG. 38

15/20




FIG. 39

16/20

ABX-EGF: Inhibition of Human Epidermoid
Carcinoma Growth in Nude Mice




FIG. 40

17/20

*Inhibition of Human Epidermoid Carcinoma
Formation in Nude Mice by ABX-EGF*

<i>Treatment</i>	<i>Dose (mg)</i>	<i>Tumor Formation^b (incidence)</i>	<i>Tumor size^c (cm³)</i>
PBS		6/6	1.376
Human IgG2 ^a	1	6/6	1.727
E7.6.3	1	0/5	0
	0.2	0/4	0
E2.5	1	0/3	0
	0.2	0/3	0
E1.1	1	0/3	0

^a control human myeloma IgG2

^b incidence determined 19 days post tumor inoculation

^c tumor size measured 19 days post tumor inoculation

FIG. 41

18/20





FIG. 42

19/20

Eradication of Established Human Epidermoid
Tumor in Nude Mice by ABX-EGF (E7.6.3)



A431 (5×10^6 /mouse) were injected at day 0

FIG. 43

20/20

Eradication of Established Human Epidermoid Tumor in Nude Mice by ABX-EGF (E2.5)





FIG. 44



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 16/28, A61K 39/395		A3	(11) International Publication Number: WO 98/50433
			(43) International Publication Date: 12 November 1998 (12.11.98)
(21) International Application Number: PCT/US98/09160		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 5 May 1998 (05.05.98)			
(30) Priority Data: 08/851,362 5 May 1997 (05.05.97) US			
(71) Applicant: ABGENIX, INC. [US/US]; 7601 Dumbarton Circle, Fremont, CA 94555 (US).			
(72) Inventors: JAKOBOVITS, Aya; 2021 Monterey Avenue, Menlo Park, CA 94025 (US). YANG, Xiao-Dong; 2833 Bryant Street, Palo Alto, CA 94306 (US). GALLO, Michael; 1376 Klamath Drive, San Jose, CA 95130 (US). JIA, Xiao-Chi; 64 Burbank Avenue, San Mateo, CA 94403 (US).		Published With international search report.	
(74) Agents: HALEY, James, F. et al.; Fish & Neave, 1251 Avenue of the Americas, New York, NY 10020 (US).		(88) Date of publication of the international search report: 4 February 1999 (04.02.99)	

(54) Title: MAN MONOCLONAL ANTIBODIES TO EPIDERMAL GROWTH FACTOR RECEPTOR



(57) Abstract

In accordance with the present invention, there are provided fully human monoclonal antibodies against human epidermal growth factor receptor (EGF-r). Nucleotide sequences encoding and amino acid sequences comprising heavy and light chain immunoglobulin molecules, particularly sequences corresponding to contiguous heavy and light chain sequences from CDR1 through CDR3, are provided. Hybridomas expressing such immunoglobulin molecules and monoclonal antibodies are also provided.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/09160

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07K16/28 A61K39/395

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	M. MENDEZ ET AL.: "Functional transplant of megabase human immunoglobulin loci recapitulates human antibody response in mice." NATURE GENETICS, vol. 15, February 1997, pages 146-156, XP002067603 New York, NY, USA see abstract see page 151, right-hand column, line 6 - line 27 see figures 5B, 6B ----- -/-	1-46

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- ✓ document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- 5 document member of the same patent family

Date of the actual completion of the international search

16 November 1998

Date of mailing of the international search report

01/12/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Nootj, F

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/09160

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	J. CORVALAN ET AL.: "Generation of fully human high affinity monoclonal antibodies to EGF receptor in mice." THE JOURNAL OF ALLERGY AND CLINICAL IMMUNOLOGY, vol. 99, no. 1 part 2, suppl. 's!', 1 January 1997, page s15 XP002067605 St. Louis, MO, USA see abstract 869	1-46
A	--- A. JAKOBOVITS ET AL.: "Humoral immunity in mice engineered with megabase human heavy and kappa light chain YACs." THE JOURNAL OF ALLERGY AND CLINICAL IMMUNOLOGY, vol. 99, no. 1 part 2, suppl. 's!', 1 January 1997, page s113 XP002067604 St. Louis, MO, USA see abstract 460	1-46
A	--- T. MATSUMOTO ET AL.: "Chimeric monoclonal antibody (Mab) C225 to the epidermal growth factor receptor (EGF-R) inhibits the growth of human transitional cell carcinoma (TCC) in the bladder of nude mice." PROCEEDINGS OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH, vol. 38, March 1997, page 229 XP002084507 USA see abstract #1543	1-46
A	--- WO 96 40210 A (IMCLONE SYSTEMS INC. & MRC COLLABORATIVE CENTRE) 19 December 1996 see examples see claims	1-46
A	--- WO 96 34096 A (CELL GENESYS INC.) 31 October 1996 see examples see claims	1-46
P,X	--- A. JAKOBOVITS: "Production and selection of antige-specific fully human monoclonal antibodies from mice engineered with human Ig loci." ADVANCED DRUG DELIVERY REVIEWS, vol. 31, no. 1-2, 6 April 1998, pages 33-42, XP002084508 Amsterdam, The Netherlands see page 39, left-hand column, line 34 - page 40, left-hand column, line 22 --- -/--	1-46

INTERNATIONAL SEARCH REPORT

Int .tional Application No

PCT/US 98/09160

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>A. JAKOBOVITS: "The long-awaited magic bullets: therapeutic human monoclonal antibodies from transgenic mice." EXPERT OPINION ON INVESTIGATIONAL DRUGS, vol. 7, no. 4, April 1998, pages 607-614, XP002084509 London, gb see page 612, left-hand column, line 10 - right-hand column, line 27 see table 1 see figures 1,3</p>	1-46
T	<p>WO 98 24893 A (ABGENIX, INC.) 11 June 1998 see the whole document</p>	1-46

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/09160

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim 46
is directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/09160


Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9640210 A	19-12-1996	AU 6267896 A CA 2222231 A EP 0831880 A	30-12-1996 19-12-1996 01-04-1998
WO 9634096 A	31-10-1996	AU 2466895 A EP 0823941 A	18-11-1996 18-02-1998
WO 9824893 A	11-06-1998	AU 5702298 A	29-06-1998



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07K 16/28, A61K 39/395		A3	(11) International Publication Number: WO 98/50433
			(43) International Publication Date: 12 November 1998 (12.11.98)
(21) International Application Number: PCT/US98/09160		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 5 May 1998 (05.05.98)		<p>Published <i>With international search report.</i></p> <p>(88) Date of publication of the international search report: 4 February 1999 (04.02.99)</p>	
(30) Priority Data: 08/851,362 5 May 1997 (05.05.97) US			
(71) Applicant: ABGENIX, INC. [US/US]; 7601 Dumbarton Circle, Fremont, CA 94555 (US).			
(72) Inventors: JAKOBOVITS, Aya; 2021 Monterey Avenue, Menlo Park, CA 94025 (US). YANG, Xiao-Dong; 2833 Bryant Street, Palo Alto, CA 94306 (US). GALLO, Michael; 1376 Klamath Drive, San Jose, CA 95130 (US). JIA, Xiao-Chi; 64 Burbank Avenue, San Mateo, CA 94403 (US).			
(74) Agents: HALEY, James, F. et al.; Fish & Neave, 1251 Avenue of the Americas, New York, NY 10020 (US).			

(54) Title: HUMAN MONOCLONAL ANTIBODIES TO EPIDERMAL GROWTH FACTOR RECEPTOR



(57) Abstract

In accordance with the present invention, there are provided fully human monoclonal antibodies against human epidermal growth factor receptor (EGF-r). Nucleotide sequences encoding and amino acid sequences comprising heavy and light chain immunoglobulin molecules, particularly sequences corresponding to contiguous heavy and light chain sequences from CDR1 through CDR3, are provided. Hybridomas expressing such immunoglobulin molecules and monoclonal antibodies are also provided.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		